Abundance for Kaehler threefolds

Frédéric Campana, Andreas Hoering, Thomas Peternell

► To cite this version:

Frédéric Campana, Andreas Hoering, Thomas Peternell. Abundance for Kaehler threefolds. 2014. hal-00958609v1

HAL Id: hal-00958609

<https://hal.science/hal-00958609v1>

Preprint submitted on 12 Mar 2014 (v1), last revised 20 Apr 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ABUNDANCE FOR KÄHLER THREEFOLDS

FRÉDÉRIC CAMPANA, ANDREAS HÖRING, AND THOMAS PETERNELL

ABSTRACT. Let X be a compact Kähler threefold with terminal singularities such that K_X is nef. We prove that K_X is semiample.

1. INTRODUCTION

1.A. Main results. Since the 1990's, the minimal model program for smooth projective threefolds is complete: every such manifold X admits a birational model X' , which is $\mathbb{Q}$ -factorial with only terminal singularities such that X' either carries a Fano fibration, in particular is uniruled, or the canonical bundle $K_{X'}$ is semiample, i.e., some positive multiple $mK_{X'}$ is generated by global sections. There are basically two parts in the program: first to establish the existence of a model X' which is either a Mori fibre space or has nef canonical divisor, and then to show that nefness implies semi-ampleness. This second part, known as “abundance”, is established by [Miy87, Kaw92a, Kwc92].

The aim of the present paper is to fully establish the minimal model program in the category of Kähler threefolds. The first part of the program, i.e., the existence of a bimeromorphic model X' which is either a Mori fibre space or has nef canonical divisor was carried out in the papers [HP13a] and [HP13b]. Thus it remains to show that nefness of the canonical divisor implies semi-ampleness, i.e. abundance holds for Kähler threefolds:

1.1. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. Then K_X is semi-ample, that is some positive multiple mK_X is globally generated.*

The paper [DP03] established the existence of some section in mK_X for non-algebraic minimal models, so the assumption above implies $\kappa(X) \geq 0$. In [Pet01] abundance was shown for non-algebraic minimal models, excluding however the case when X has no non-constant meromorphic function. Our arguments do not use any information on the structure of X and work both for algebraic and non-algebraic Kähler threefolds.

As a corollary, we establish a longstanding conjecture on Kähler threefolds:

1.2. Theorem. *Let X be smooth compact Kähler threefold. Assume that X is simple, i.e. there is no positive-dimensional proper subvariety through the very general point of X . Then there exists a bimeromorphic morphism $X \rightarrow T/G$ where T is a torus and G a finite group acting on T .*

Date: March 12, 2014.

2000 Mathematics Subject Classification. 32J27, 14E30, 14J30, 32J17, 32J25.

Key words and phrases. MMP, rational curves, Zariski decomposition, Kähler manifolds, abundance.

It was known since some time that this result is a consequence of the existence of good minimal models [Pet98, p.731]; for the convenience of the reader, we give the proof at the end of the paper.

1.B. Outline of the paper. In principle, we follow the strategy of [Kaw92a], [Kwc92] to settle the abundance theorem in the Kähler case. More precisely, consider a normal compact $\mathbb{Q}$ -factorial Kähler threefold X with only terminal singularities and nef anticanonical divisor K_X . Denote by $\kappa(X)$ its Kodaira dimension and by $\nu(X)$ the numerical dimension, which, is defined as

$$\nu(X) := \max\{m \in \mathbb{N} \mid c_1(K_X)^m \neq 0\}.$$

Both invariants are subject to the inequality $\kappa(X) \leq \nu(X)$, with equality if K_X is semi-ample. Conversely, as Kawamata observed in [Kaw85, Thm.6.1], in order to prove abundance, it is sufficient to prove equality: $\kappa(X) = \nu(X)$. By the base-point free theorem and an argument of Kawamata [Kaw85, Thm.7.3] the main challenge is to rule out the possibility

$$\kappa(X) = 0 \text{ and } 0 < \nu(X) < 3.$$

Since we know that $\kappa(X) \geq 0$, there exists a positive number m and an effective divisor D such that $D \in |mK_X|$. Since D might be very singular it is however not possible to analyse the divisor D directly. In order to circumvent this difficulty, Kawamata developed a strategy, further explored in [Kwc92], to improve the situation via certain birational transformations. This requires deep techniques of birational geometry of pairs within the theory of minimal models. In particular we have to run a log MMP for certain lc pairs (X, Δ) .

Therefore the first part of the paper (Sections 3 and 4) establishes the relevant results. Although we are not able to prove the cone and contraction theorems for arbitrary lc (or dlt) pairs, the additional conditions on the boundary Δ and the singularities of X will be satisfied in the applications we are interested in.

1.3. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with rational singularities that is not uniruled. Let Δ be an effective $\mathbb{Q}$ -divisor on X such that the pair (X, Δ) is lc. Then there exists an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X and a number d such that*

$$0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$$

and

$$\overline{NA}(X) = \overline{NA}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i]$$

If (X, Δ) is dlt, then we may take the curves Γ_i to be rational and $d = 4$.

The dual Kähler cone $\overline{NA}(X)$ replaces the Mori cone of curves $\overline{NE}(X)$ which is obviously too small in the non-algebraic setting (cf. [HP13a, Sect.1]). We will prove Theorem 1.3 on page 21 as a consequence of the weak cone theorem 4.2 and some partial contraction results. As in the projective case [Fuj11a], it is very likely that the rays can be represented by rational curves with bound $d = 4$.

1.4. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold that is not uniruled. Let Δ be an effective $\mathbb{Q}$ -divisor such that the pair (X, Δ) is dlt. Let R be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$. If the ray R is divisorial with exceptional divisor S , suppose that S has slc singularities.*

Then the contraction of R exists in the Kähler category.

Based on these results we can follow the arguments of [Kwc92, Ch.13, 14] to replace the threefold X with some bimeromorphic model having a pluricanonical divisor $D \in |mK_X|$ such that D_{red} is not too singular (cf. Lemmata 6.1, 6.6, 6.7, 6.9). With this preparation one can exclude the case $\kappa(X) = 0, \nu(X) = 1$ following a deformation argument of Miyaoka [Miy88] [Kwc92, Ch.11]. For the case $\nu(X) = 2$ the idea is to prove via a Riemann-Roch computation that $h^0(X, mK_X)$ grows linearly. In the literature [Kaw92a], [Kwc92, Ch.14] this computation is based on the Chern class inequality

$$(K_X + B) \cdot \hat{c}_2(X) \geq 0$$

where (X, B) is an lc pair such that $K_X + B$ is nef and $\hat{c}_2(X)$ the second Chern class of the $\mathbb{Q}$ -sheaf $\hat{\Omega}_X$ (cf. [Kwc92, Ch.10]). Since an lc pair is not necessarily smooth in codimension one the proof of this inequality is quite involved. We simplify this argument as follows: let $X' \rightarrow X$ be a terminal modification of the pair (X, B) . Since X' is smooth in codimension one the Riemann-Roch computation can be done using the classical second Chern class $c_2(X')$. Although the (log-)canonical divisor of X' is not necessarily nef, a generalisation of Miyaoka's inequality for the second Chern class (Theorem 7.11) allows us to conclude. We think that this new strategy might be of independent interest.

Acknowledgements. We thank the Forschergruppe 790 "Classification of algebraic surfaces and compact complex manifolds" of the Deutsche Forschungsgemeinschaft for financial support. A. Höring was partially also supported by the A.N.R. project CLASS¹.

2. NOTATION AND BASIC FACTS

We will use frequently standard terminology of the minimal model program (MMP) as explained in [KM98] or [Deb01]. We will also use without further comment results which are stated for algebraic varieties in [KM98], [Kwc92] if their proof obviously works for complex spaces.

Recall that a normal complex space X is $\mathbb{Q}$ -factorial if for every Weil divisor D there exists an integer $m \in \mathbb{N}$ such that $\mathcal{O}_X(mD)$ is a locally free sheaf, i.e. mD is a Cartier divisor. Since however the canonical sheaf $K_X = \omega_X$ need not be a $\mathbb{Q}$ -Weil divisor, we include in the definition of $\mathbb{Q}$ -factoriality also the condition that there is a number $m \in \mathbb{N}$ such that the coherent sheaf

$$(K_X^{\otimes m})^{**} = (\omega_X^{\otimes m})^{**}$$

is locally free. We shall write

$$mK_X = (K_X^{\otimes m})^{**}$$

for short.

2.1. Definition. Let X be a normal complex space. A boundary divisor is an effective $\mathbb{Q}$ -divisor $\Delta = \sum_i a_i \Delta_i$ such that $0 \leq a_i \leq 1$.

¹ANR-10-JCJC-0111

Given a boundary divisor Δ , we refer to (X, Δ) as a “pair”. For pairs (X, Δ) , the notions “lc” (log-canonical), “klt” (Kawamata log-terminal), “dlt” (divisorial log-terminal) and “terminal” are defined exactly as in the algebraic context; we refer to [KM98].

2.2. Definition. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler space that is not uniruled, and let Δ be a boundary divisor on X such that the pair (X, Δ) is lc (resp. dlt, klt, terminal). Set $(X_0, \Delta_0) := (X, \Delta)$. A terminating $(K + \Delta)$ -MMP is a finite sequence of bimeromorphic maps*

$$\varphi_i : (X_i, \Delta_i) \dashrightarrow (X_{i+1}, \Delta_{i+1} := (\varphi_i)_* \Delta_i)$$

where $i \in \{0, \dots, n\}$ for some $n \in \mathbb{N}$ that has the following properties:

- a) For every $i \in \{0, \dots, n+1\}$, the complex space X_i is a normal $\mathbb{Q}$ -factorial compact Kähler space and the pair (X_i, Δ_i) is lc (resp. dlt, klt, terminal).
- b) For every $i \in \{0, \dots, n\}$, the map φ_i is either the contraction of a $(K_{X_i} + \Delta_i)$ -negative extremal ray $R_i \in \overline{NA}(X_i)$ that is divisorial or the flip of a $K_{X_i} + \Delta_i$ -negative extremal ray $R_i \in \overline{NA}(X_i)$ that is small.
- c) The class $K_{X_{n+1}} + \Delta_{n+1}$ is nef.

We will abbreviate such a $(K + \Delta)$ -MMP by

$$(X, \Delta) \dashrightarrow (X', \Delta')$$

where $(X', \Delta') := (X_{n+1}, \Delta_{n+1})$ and the bimeromorphic map is the composition $\varphi_n \circ \dots \circ \varphi_0$.

In the definition, $\overline{NA}(X)$ denotes the “dual Kähler”, as defined in [HP13a, 3.8]. We will review this notion in Definition 3.4 below.

2.A. Bimeromorphic models. Bimeromorphic models arising as partial resolutions of singularities play an important role in the recent development of the MMP for projective manifolds. The existence of these models is usually - in the algebraic setting, see [Fuj11b, 10.4], [KK10, 3.1], [Kol13, Ch.1.4] - derived as a consequence of the existence and termination of some MMP for pairs, which is at this point not yet established in the Kähler category. We may nevertheless assume their existence since they are constructed by taking first a log-resolution $Y' \rightarrow X$ of the pair (X, Δ) and then running some MMP *over* X . Since we may choose the log-resolution to be a projective morphism, the usual relative versions of the cone and contraction theorem [Nak87] [KM98, Ch.3.6] together with termination results in dimension three allows to conclude. The Kähler property of Y follows again from [Var89, 1.3.1], since μ is a projective morphism. We will now apply this argument in two situations:

2.3. Theorem. *Let X be a normal compact Kähler threefold with lc singularities, i.e., $(X, 0)$ is lc. Then there exists a bimeromorphic morphism $\mu : Y \rightarrow X$ from a normal $\mathbb{Q}$ -factorial Kähler threefold Y with terminal singularities and a boundary divisor Δ_Y such that the pair (Y, Δ_Y) is lc and*

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X.$$

Proof. Let $\pi : \hat{X} \rightarrow X$ be a resolution of singularities. Let $\hat{X} \dashrightarrow Y$ be a $K_\bullet$ -MMP over X . The outcome of this MMP is a normal $\mathbb{Q}$ -factorial threefold $\mu : Y \rightarrow X$ with terminal singularities such that K_Y is μ -nef. We write

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X,$$

where the support of Δ_Y is contained in the μ -exceptional locus. By the negativity lemma [KM98, Lemma 3.39] we see that Δ_Y is effective and since X has lc singularities, Δ_Y is a boundary divisor. The pair (Y, Δ_Y) is lc since this property is invariant under crepant bimeromorphic morphisms [KM98, Lemma 2.30]. $\square$

2.4. Theorem. *Let X be a normal compact Kähler threefold, and let Δ be a boundary divisor on X such that the pair (X, Δ) is lc. Then (X, Δ) has a dlt model, i.e. there exists a bimeromorphic morphism from a normal $\mathbb{Q}$ -factorial Kähler threefold Y and a boundary divisor Δ_Y such that the pair (Y, Δ_Y) is dlt and*

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^*(K_X + \Delta).$$

Proof. The same reasons as in Theorem 2.3 applies also here; we refer to [KK10, 3.1] and [Fuj11b, 10.4], based on [BCHM10, Thm.1.2], which is a result on projective morphisms. The deduction from [BCHM10] is basically a computation of discrepancies including an application of the negativity lemma. $\square$

2.B. Adjunction. Let X be a normal complex space and let $S \subset X$ be a prime divisor that is $\mathbb{Q}$ -Cartier. Let B be an effective $\mathbb{Q}$ -divisor such that $K_X + B$ is $\mathbb{Q}$ -Cartier and $S \not\subset \text{Supp}(B)$. Let $\nu : \tilde{S} \rightarrow S$ be the normalisation, then Shokurov's differend [Sho92, Ch.3] is a naturally defined effective $\mathbb{Q}$ -divisor $B_{\tilde{S}}$ on $\tilde{S}$ such that

$$(1) \quad K_{\tilde{S}} + B_{\tilde{S}} \sim_{\mathbb{Q}} \nu^*(K_X + S + B)|_S,$$

From the construction in [Sho92, Ch.3] one sees immediately that

$$\text{Supp } \nu_*(B_{\tilde{S}}) \subset S_{\text{sing}} \cup (\text{Supp } B \cap S) \cup (X_{\text{sing}} \cap S).$$

Suppose now that $\dim X = 3$ and let $\mu : \hat{S} \rightarrow \tilde{S}$ be the minimal resolution. Then we have

$$K_{\hat{S}} \sim_{\mathbb{Q}} \mu^* K_{\tilde{S}} - N,$$

where N is an effective μ -exceptional divisor [Sak84, 4.1]. Set $E := N + \mu^* B_{\tilde{S}}$, then E is a canonically defined effective $\mathbb{Q}$ -divisor, such that

$$(2) \quad K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S + B)|_S,$$

where $\pi : \hat{S} \rightarrow S$ is the composition $\nu \circ \mu$. Since E is μ -exceptional, we obtain

$$\text{Supp } \pi_*(E) \subset S_{\text{sing}} \cup (\text{Supp } B \cap S) \cup (X_{\text{sing}} \cap S).$$

Let $C \subset S$ be a curve such that

$$C \not\subset S_{\text{sing}} \cup (\text{Supp } B \cap S) \cup (X_{\text{sing}} \cap S).$$

Then the morphism π is an isomorphism in the generic point of C , and we can define the strict transform $\hat{C} \subset \hat{S}$ as the closure of $C \setminus S_{\text{sing}}$. By what precedes, $\hat{C} \not\subset E$, thus by the projection formula and (2), we deduce

$$(3) \quad K_{\hat{S}} \cdot \hat{C} \leq (K_{\hat{S}} + E) \cdot \hat{C} = (K_X + S + B) \cdot C.$$

3. SINGULAR KÄHLER SPACES

3.A. Bott-Chern cohomology. In this section we review the cohomology groups that replace the Néron-Severi space $\text{NS}(X) \otimes \mathbb{R}$ from the projective setting. In [HP13a] we used the space

$$H_{\partial\bar{\partial}}^{1,1}(X) = \{\omega \in \mathcal{A}^{1,1}(X) \mid d\omega = 0\} / \partial\bar{\partial}\mathcal{A}^0(X)$$

to define the Néron-Severi space $N^1(X)$ as the subspace consisting of the classes of real $(1,1)$ -forms. Since all objects we are working with are actually $\partial\bar{\partial}$ -closed, it seems advisable to unwind the definition, following [BPEG13]. However, we will obtain the same cohomology space. The advantage of this new definition is that it considerably simplifies the proofs of some technical statements. We recall that $\mathcal{A}_X^{p,q}$ is the sheaf of $\mathcal{C}^\infty$ -forms of type (p,q) , and $\mathcal{A}_X^0$ is the sheaf of $\mathcal{C}^\infty$ -functions; $\mathcal{D}_X$ denotes the sheaf of distributions.

3.1. Definition. [BPEG13, Defn. 4.6.2] *Let X be a normal complex space. Let $\mathcal{H}_X$ be the sheaf of real parts of holomorphic functions multiplied² with i . A $(1,1)$ -form (resp. $(1,1)$ -current) with local potentials on X is a global section of the quotient sheaf $\mathcal{A}_X^0/\mathcal{H}_X$ (resp. $\mathcal{D}_X/\mathcal{H}_X$). We define the Bott-Chern cohomology*

$$H_{\text{BC}}^{1,1}(X) := H^1(X, \mathcal{H}_X).$$

Using the exact sequence

$$0 \rightarrow \mathcal{H}_X \rightarrow \mathcal{A}_X^0 \rightarrow \mathcal{A}_X^0/\mathcal{H}_X \rightarrow 0,$$

and the fact that $\mathcal{A}_X^0$ is acyclic, we obtain a surjective map

$$H^0(X, \mathcal{A}_X^0/\mathcal{H}_X) \rightarrow H^1(X, \mathcal{H}_X).$$

Thus we can see an element of the Bott-Chern cohomology group as a closed $(1,1)$ -form with local potentials modulo all the forms that are globally of the form $dd^c u$. Using the exact sequence

$$0 \rightarrow \mathcal{H}_X \rightarrow \mathcal{D}_X \rightarrow \mathcal{D}_X/\mathcal{H}_X \rightarrow 0,$$

we see that one obtains the same Bott-Chern group, if we consider $(1,1)$ -currents with local potentials.

A fundamental property (shown in the proof of [BPEG13, Lemma 4.6.1]) is that for any proper bimeromorphic morphism $\pi : X \rightarrow Y$ between normal complex spaces we have

$$(4) \quad \pi_*(\mathcal{H}_X) = \mathcal{H}_Y.$$

Suppose now that X is a compact normal space in the Fujiki class $\mathcal{C}$ with at most rational singularities. Consider the exact sequence

$$(5) \quad 0 \rightarrow \mathbb{R} \rightarrow \mathcal{O}_X \rightarrow \mathcal{H}_X \rightarrow 0,$$

then the long exact sequence in cohomology yields

$$0 \rightarrow H^1(X, \mathbb{R}) \rightarrow H^1(X, \mathcal{O}_X) \rightarrow H^1(X, \mathcal{H}_X) \rightarrow H^2(X, \mathbb{R}) \rightarrow \dots$$

²We “twist” the definition from [BPEG13] in order to get a group that injects in $H^2(X, \mathbb{R})$ rather than $H^2(X, i\mathbb{R})$

Since $H^1(X, \mathbb{R}) \rightarrow H^1(X, \mathcal{O}_X)$ is an isomorphism³, we obtain an injection

$$(6) \quad H^1(X, \mathcal{H}_X) \hookrightarrow H^2(X, \mathbb{R}).$$

Let now $\varphi : X \rightarrow Y$ be a proper bimeromorphic morphism between compact normal spaces in class $\mathcal{C}$ with at most rational singularities. Using (4), the Leray spectral sequence yields an exact sequence

$$0 \rightarrow H^1(Y, \mathcal{H}_Y) \rightarrow H^1(X, \mathcal{H}_X) \rightarrow H^0(Y, R^1\varphi_*(\mathcal{H}_X)).$$

In particular we have an injection between the Bott-Chern spaces (cp. [HP13a, Lemma 3.3])

$$(7) \quad H_{\text{BC}}^{1,1}(Y) \rightarrow H_{\text{BC}}^{1,1}(X).$$

Since Y has only rational singularities and φ is bimeromorphic, the Grothendieck spectral sequence immediately yields

$$R^q\varphi_*(\mathcal{O}_X) = 0$$

for $q \geq 1$. Consequently $R^1\varphi_*(\mathbb{R}) = 0$, so the Leray spectral sequence yields an exact sequence

$$0 \rightarrow H^2(Y, \mathbb{R}) \rightarrow H^2(X, \mathbb{R}) \rightarrow H^0(Y, R^2\varphi_*(\mathbb{R})).$$

Moreover the push-forward of the exact sequence (5) yields an isomorphism

$$R^1\varphi_*(\mathcal{H}_X) \simeq R^2\varphi_*(\mathbb{R}).$$

Combining this isomorphism and the sequences above with the injection (6) we obtain a commutative diagram

$$(8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^1(Y, \mathcal{H}_Y) & \longrightarrow & H^1(X, \mathcal{H}_X) & \longrightarrow & H^0(Y, R^1\varphi_*(\mathcal{H}_X)) \\ & & \downarrow & & \downarrow & & \downarrow \simeq \\ 0 & \longrightarrow & H^2(Y, \mathbb{R}) & \longrightarrow & H^2(X, \mathbb{R}) & \longrightarrow & H^0(Y, R^2\varphi_*(\mathbb{R})) \end{array}$$

We will now use the claim (*) from the proof of [KM92, Thm.12.1.3]: given an element $S \in H^2(X, \mathbb{R})$ such that its image in $H^0(Y, R^2\varphi_*(\mathbb{R}))$ is non-zero, there exists an element $C \in H_2(X/Y, \mathbb{R})$ such that $S \cdot C \neq 0$. Since $H_2(X/Y, \mathbb{R})$ is generated by classes of algebraic cycles [KM92, Thm.12.1.3] we obtain the following (cp. [HP13a, Lemma 8.4]).

3.2. Lemma. *Let $\varphi : X \rightarrow Y$ be a proper bimeromorphic morphism between compact normal spaces in class $\mathcal{C}$ with at most rational singularities. Then we have an injection*

$$\varphi^* : H^1(Y, \mathcal{H}_Y) \hookrightarrow H^1(X, \mathcal{H}_X)$$

and

$$\text{Im } \varphi^* = \{\alpha \in H^1(X, \mathcal{H}_X) \mid \alpha \cdot C = 0 \quad \forall C \subset X \text{ curve s.t. } \varphi(C) = pt\}.$$

Furthermore, let $\alpha \in H^1(X, \mathcal{H}_X) \subset H^2(X, \mathbb{R})$ be a class such that $\alpha = \varphi^*\beta$ with $\beta \in H^2(Y, \mathbb{R})$. Then there exists a smooth real closed $(1,1)$ -form ω_Y on Y such that $\alpha = \varphi^*[\omega_Y]$.

³Since X has rational singularities we have $H^1(\hat{X}, \mathbb{R}) \simeq H^1(X, \mathbb{R})$ and $H^1(\hat{X}, \mathcal{O}_{\hat{X}}) \simeq H^1(X, \mathcal{O}_X)$ where $\hat{X} \rightarrow X$ is a resolution by a compact Kähler manifold. Now we apply the classical result from the Kähler case.

Proof. A simple diagram chase in (8). $\square$

At this point we explain the connection to [HP13a]:

3.3. Proposition. *Let X be a normal compact space in class $\mathcal{C}$ with at most rational singularities. Then*

$$H^1(X, \mathcal{H}_X) = H_{\text{BC}}^{1,1}(X) \simeq N^1(X).$$

Proof. Obviously, we have a canonical injective map

$$H_{\text{BC}}^{1,1}(X) \rightarrow N^1(X).$$

Choose a desingularisation $\pi : \hat{X} \rightarrow X$ with $\hat{X}$ Kähler. Then the pull-back morphism $H_{\text{BC}}^{1,1}(X) \rightarrow H_{\text{BC}}^{1,1}(\hat{X})$ (resp. $N^1(X) \rightarrow N^1(\hat{X})$) is injective by (7) (resp. [HP13a, Lemma 3.3]). Since $H_{\text{BC}}^{1,1}(\hat{X}) \simeq N^1(\hat{X})$ things come down to prove the following: for every

$$\alpha \in H_{\text{BC}}^{1,1}(\hat{X}) \cap \pi^*(N^1(X))$$

we have $\alpha \in \pi^*(H_{\text{BC}}^{1,1}(X))$. However $\alpha \in \pi^*(H^2(X, \mathbb{R}))$ by the second line of (8), consequently $\alpha \in \pi^*(H_{\text{BC}}^{1,1}(X))$ by Lemma 3.2. $\square$

3.4. Definition. *Let X be a normal compact complex space in class $\mathcal{C}$. We define $N_1(X)$ to be the vector space of real closed currents of bidimension $(1, 1)$ modulo the following equivalence relation: $T_1 \equiv T_2$ if and only if*

$$T_1(\eta) = T_2(\eta)$$

for all real closed $(1, 1)$ -forms η (with local potentials).

We define $\overline{NA}(X) \subset N_1(X)$ to be the closed cone generated by the classes of positive closed currents. The closed cone of curves is the subcone

$$\overline{NE}(X) \subset \overline{NA}(X)$$

of those positive closed currents arising as currents of integration over curves.

3.5. Remark. In Definition it is irrelevant whether the forms η have local potentials or not. In fact, in both cases the space $N_1(X)$ is dual to $N^1(X)$; see Proposition 3.7 in [HP13a]. The proof of this proposition also works in the case of forms with local potentials (using the obvious fact that a closed current T of bidimension T .

The closedness of a current of integration T_C with C an irreducible curves follows by passing to a desingularisation π and using the closeness of the current $T_{\hat{C}}$, where $\hat{C}$ is the strict transform, if C is not contained in the singular locus of X . The integral $\int_C \phi$ for a form ϕ on X is well-defined, even if $C \subset \text{Sing}(X)$. In this case we take $\hat{C}$ to be any curve $\hat{C}$ such that $\pi(\hat{C}) = C$ and observe that $\pi_*(T_{\hat{C}}) = dT_C$, where d is the degree of $\pi|_{\hat{C}}$. Hence T_C is closed.

For the proof of the contraction theorem 1.4 we will need the following statements:

3.6. Proposition. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler space. Let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$. Suppose that there exists a bimeromorphic morphism $\varphi : X \rightarrow Y$ onto a normal complex space Y such that $-(K_X + \Delta)$ is φ -ample and a curve $C \subset X$ is contracted if and only if $[C] \in \mathbb{R}^+[\Gamma_i]$.*

a) Then we have two exact sequences

$$(9) \quad 0 \rightarrow H^2(Y, \mathbb{R}) \xrightarrow{\varphi^*} H^2(X, \mathbb{R}) \xrightarrow{\alpha \mapsto \alpha \cdot \Gamma_i} \mathbb{R} \rightarrow 0$$

$$(10) \quad 0 \rightarrow N^1(Y) \xrightarrow{\varphi^*} N^1(X) \xrightarrow{[L] \mapsto L \cdot \Gamma_i} \mathbb{R} \rightarrow 0.$$

In particular we have $b_2(X) = b_2(Y) + 1$.

b) We have an exact sequence

$$(11) \quad 0 \rightarrow \text{Pic}(Y) \xrightarrow{\varphi^*} \text{Pic}(X) \xrightarrow{[L] \mapsto L \cdot \Gamma_i} \mathbb{Z}.$$

- c) If the contraction is divisorial, the variety Y is $\mathbb{Q}$ -factorial and its Picard number is $\rho(X) - 1$. Moreover the pair $(Y, \varphi_*\Delta)$ is dlt
- d) If the contraction is small with flip $\varphi^+ : X^+ \rightarrow Y$, the complex space X^+ is $\mathbb{Q}$ -factorial and its Picard number is $\rho(X)$. Moreover the pair $(X^+, (\varphi^+)^{-1}_* \circ \varphi_*\Delta)$ is dlt.

Proof. Since $-(K_X + \Delta)$ is φ -ample, the morphism φ is projective. Since (X, Δ) is dlt and X is $\mathbb{Q}$ -factorial, the pair $(X, (1 - \varepsilon)\Delta)$ is klt for small positive ε . Since $-(K_X + (1 - \varepsilon)\Delta)$ is φ -ample, possibly after passing to a smaller ε , we obtain $R^q\varphi_*(\mathcal{O}_X) = 0$ for $q \geq 1$ by [Anc87, Thm.3.1]. Since X has rational singularities ([KM98, 5.22]), so does Y ; moreover a) is proved by (8). The property b) follows from [KM98, Thm.3.25(4)] and implies the properties c) and d) as in [KM98, Prop.3.36, Prop.3.37, Cor.3.44]. $\square$

3.7. Lemma. Let X be a normal $\mathbb{Q}$ -factorial compact Kähler space, and let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let

$$\varphi : X \rightarrow Y \text{ resp. } \varphi : X \dashrightarrow X^+$$

be the divisorial contraction resp. the flip of a $K_X + \Delta$ -negative extremal ray $R \in \overline{NA}(X)$. Then the following holds:

- a) Let D be a nef divisor such that $D \cdot R = 0$. Then $\varphi_*(D)$ is nef and

$$\nu(D) = \nu(\varphi_*(D)).$$

Moreover if $S \subset X$ is a prime divisor that is not contracted by φ , then $D \cdot S \neq 0$ in $H^4(X, \mathbb{R})$ if and only if $\varphi_*(D) \cdot \varphi(S) \neq 0$ in $H^4(Y, \mathbb{R})$.

- b) Let B be a boundary divisor such that the pair (X, B) is lc. If $(K_X + B) \cdot R = 0$, then the pair $(Y, \varphi_*(B))$ is lc.

Proof. Set $L := \mathcal{O}_X(D)$. We first treat the case when φ is divisorial: by Proposition 3.6,b) we have $L \simeq \varphi^*M$ with M a line bundle on Y and obviously $M \simeq \mathcal{O}_Y(\varphi_*D)$. Being nef is invariant under pull-back [Pau98], so M is nef. The numerical dimension is invariant since the pull-back commutes with the intersection product. If $S' = \varphi(S)$, then $L|_S = (\varphi|_S)^*(M|_{S'})$, hence $L \cdot S \neq 0$ if and only if $M \cdot \varphi(S) \neq 0$ by the projection formula. For the property b) we refer to [KM98, Lemma 2.30].

In the flipping case, consider the small contraction $f : X \rightarrow Y$ associated to R and the flip $f^+ : X^+ \rightarrow Y$. For a) apply the considerations above first to f and then to f^+ . For b), apply [KM98, 2.30] to see that $(Y, f_*(B))$ is lc and then again to conclude that $(X^+, \varphi_*(B) = (f^+)^{-1}_* \circ f_*(B))$ is lc. $\square$

3.B. Kähler criteria. In this subsection we generalize some Kähler criteria given in [HP13a] to threefolds with non-isolated singularities.

3.8. Theorem. *Let X be a normal compact threefold in class $\mathcal{C}$. Let $\eta \in \mathcal{A}^{1,1}(X)$ be a closed real $(1,1)$ -form such that $T(\eta) > 0$ for all $[T] \in \overline{NA}(X) \setminus 0$. Suppose that for every irreducible curve $C \subset X$ we have $[C] \neq 0$ in $N_1(X)$. Then $\{\eta\} \in N^1(X)$ is represented by a Kähler class, in particular X is Kähler.*

Proof. We slightly generalise the arguments of [HP13a, Thm.3.16] by removing the assumption that the singularities are isolated. If X is smooth, then [HP13a, Thm.3.16] applies, so that $\{\eta\}$ is a Kähler class.

We shall now reduce ourselves to this case by considering a resolution $\mu : \hat{X} \rightarrow X$ with $\hat{X}$ a compact Kähler manifold such that a suitable exceptional divisor E of $\hat{X}$ is μ -ample. As in [HP13a], we argue that the class of an irreducible curve in $\hat{X}$ does not vanish. Furthermore, we check as in Step 2 of the proof of [HP13a, Thm.3.16] that

$$T(\mu^*(\eta) - \frac{1}{n}E) > 0$$

for all classes $[T] \in \overline{NA}(\hat{X})$ if n is a sufficiently big positive integer. Hence, by the solution in the smooth case, we may choose a Kähler form $\hat{\eta}$ on $\hat{X}$ in the class $\{\mu^*(\eta) - \frac{1}{n}E\}$. Consider the Kähler current

$$\{\mu_*(\hat{\eta})\} = \{\eta\}.$$

Let Z be an irreducible component of the Lelong level sets of $\mu_*(\hat{\eta})$; then Z is either a point or an irreducible curve. By our assumption, $\{\eta\}|_Z$ is a Kähler class on Z . By [DP04, Prop.0.6] this implies that $\{\eta\}$ is a Kähler class on X . $\square$

3.9. Corollary. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold that is not uniruled. Let Δ be an effective $\mathbb{Q}$ -divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$.*

Suppose that there exists a bimeromorphic morphism $\varphi : X \rightarrow Y$ such that $-(K_X + \Delta)$ is φ -ample and a curve $C \subset X$ is contracted if and only if $[C] \in \mathbb{R}^+[\Gamma_i]$. Then Y is a Kähler space.

Proof. As in the proof of Proposition 3.6 we see that Y has rational singularities. By assumption, $\mathbb{R}^+[\Gamma_i]$ is extremal in $\overline{NA}(X)$; denote by $\alpha \in N^1(X)$ a nef supporting class (cf. Proposition 4.10). Consequently

$$\alpha \cdot Z' > 0$$

for every class $Z' \in \overline{NA}(X) \setminus \mathbb{R}_0^+[\Gamma_i]$. By Proposition 3.6,a) there exists a $\beta \in N^1(Y)$ with $\alpha = \varphi^*(\beta)$ and by what precedes we have

$$\beta \cdot Z > 0$$

for any $Z \in \overline{NA}(Y) \setminus 0$.

We claim that for any irreducible curve $C \subset Y$, there exists a curve $C' \subset X$ such that $C = \varphi(C')$. Since C' is not contracted by φ we have $\alpha \cdot C' > 0$, so $\beta \cdot C > 0$ by the projection formula. In particular, $[C] \neq 0$ in $N_1(X)$, so the statement follows from Theorem 3.8.

For the proof of the claim note first that if C is not contained in the image of the exceptional locus, then we can just take the strict transform $C' \subset X$. If C is in the

image of the exceptional locus, then φ is divisorial and maps a surface S onto C . Since the morphism $\varphi|_S : S \rightarrow C$ is projective, the surface S is projective. Thus some multisection $C' \subset S$ has the required property. $\square$

4. MMP FOR PAIRS

In this section we prove several results on the MMP for pairs for Kähler threefolds. While we are not able to establish the log-MMP in full generality, the results are sufficient for the application to the abundance problem in Section 6. We start by proving the cone theorem and (parts of) the contraction theorem in the Subsections 4.A and 4.B, while Subsection 4.C merely collects the relevant results on existence of flips and termination. In order to simplify the statements we make the following

4.1. Assumption. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with rational singularities. Let $\Delta = \sum_i a_i \Delta_i$ be an effective $\mathbb{Q}$ -divisor on X such that the pair (X, Δ) is lc. We also assume that X is not uniruled, so by [HP13a, Cor.1.4] the Kodaira dimension $\kappa(\hat{X})$ of a desingularisation $\hat{X}$ is non-negative, hence $\kappa(X) \geq 0$. Thus we have*

$$(12) \quad K_X \sim_{\mathbb{Q}} \sum \lambda_j S_j,$$

where the S_j are integral surfaces in X , the coefficients $\lambda_j \in \mathbb{Q}^+$.

4.A. Cone theorem. In this section we will prove a weak form of the cone theorem for non-uniruled lc pairs. The stronger cone theorem 1.3 will then be a consequence of the contraction results in Subsection 4.B which are based on the weak cone theorem:

4.2. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with rational singularities. Let Δ be a boundary divisor on X such that the pair (X, Δ) is lc. If X is not uniruled, there exists a positive integer $d \in \mathbb{N}$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that*

$$0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$$

and

$$\overline{NA}(X) = \overline{NA}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

We will prove this statement on page 16, but this will take some technical preparation.

4.3. Lemma. *Under the Assumption 4.1, let $S \subset X$ be a prime divisor such that $(K_X + \Delta)|_S$ is not pseudoeffective. Then S is Moishezon and any desingularisation $\hat{S}$ is a uniruled projective surface.*

Proof. Let $\pi : \hat{S} \rightarrow S$ be the composition of the normalisation followed by the minimal resolution of the normalised surface. Our goal is to show that there exists an effective $\mathbb{Q}$ -divisor E such that $K_{\hat{S}} + E$ is not pseudoeffective, in particular we have $\kappa(\hat{S}) = -\infty$. It then follows from the Castelnuovo-Kodaira classification that $\hat{S}$ is covered by rational curves, in particular it is a projective surface [BHPVdV04]. Thus S is Moishezon.

1st case. Suppose that $K_X|_S$ is not pseudoeffective. Since K_X is $\mathbb{Q}$ -effective this implies that S is one of the surfaces appearing in the decomposition (12). Up to renumbering we may suppose that $S = S_1$. Observe

$$S = S_1 = \frac{1}{\lambda_1} K_X - \sum_{j=2}^r \frac{\lambda_j}{\lambda_1} S_j.$$

Applying (2) with $B = 0$, there exists an effective $\mathbb{Q}$ -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S)|_S = \frac{(\lambda_1 + 1)}{\lambda_1} \pi^* K_X|_S - \sum_{j=2}^r \frac{\lambda_j}{\lambda_1} \pi^* S_j|_S.$$

By assumption $K_X|_S$ is not pseudoeffective, and $-\sum_{j=2}^r \frac{\lambda_j}{\lambda_1} \pi^* S_j|_S$ is anti-effective. Thus $K_{\hat{S}} + E$ is not pseudoeffective.

2nd case. Suppose that $K_X|_S$ is pseudoeffective. Then $S \subset \Delta$, hence, up to renumbering, we may suppose that $S = \Delta_1$. We claim that

$$(K_X + \Delta_1 + \sum_{i \geq 2} a_i \Delta_i)|_S$$

is not pseudoeffective. Otherwise, by $0 \leq a_1 \leq 1$ and the pseudo-effectivity of $K_X|_S$, the divisor

$$(K_X + \Delta)|_S = a_1(K_X + \Delta_1 + \sum_{i \geq 2} a_i \Delta_i)|_S + (1 - a_1)(K_X + \sum_{i \geq 2} a_i \Delta_i)|_S$$

is a convex combination of pseudoeffective classes, hence itself pseudoeffective, a contradiction. Applying (2) with $S = \Delta_1$ and $B = \sum_{i \geq 2} a_i \Delta_i$, there exists an effective $\mathbb{Q}$ -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + \Delta + \sum_{i \geq 2} a_i \Delta_i)|_S.$$

Thus $K_{\hat{S}} + E$ is not pseudoeffective. $\square$

4.4. Corollary. *Under the Assumption 4.1, the divisor $K_X + \Delta$ is nef if and only if*

$$(K_X + \Delta) \cdot C \geq 0$$

for every curve $C \subset X$.

Proof. One implication is trivial. Suppose now that $K_X + \Delta$ is nef on all curves C . We will argue by contradiction and suppose that $K_X + \Delta$ is not nef. Since $K_X + \Delta$ is pseudo-effective and the restriction to every curve is nef, there exists by [Bou04, Prop.3.4] an irreducible surface $S \subset X$ such that $(K_X + \Delta)|_S$ is not pseudo-effective. By Lemma 4.3 a desingularisation $\pi : \hat{S} \rightarrow S$ of the surface S is projective, so (by [BDPP13]) there exists a covering family of curves $C_t \subset S$ such that for the strict transforms

$$\pi^*(K_X + \Delta)|_S \cdot \hat{C}_t < 0.$$

Hence we obtain $(K_X + \Delta) \cdot C_t < 0$, a contradiction. $\square$

If X is a projective manifold, the cone theorem is a consequence of Mori's bend-and-break technique. We will now show that an analogue of this technique is available for threefolds that are lc pairs.

4.5. Definition. Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold, and let Δ be a boundary divisor. We say that $K_X + \Delta$ has the bend-and-break property if there exists a positive number $d = d_{K_X + \Delta} \in \mathbb{Q}^+$ such that the following holds: given any curve $C \subset X$ such that

$$-(K_X + \Delta) \cdot C > d,$$

there exist non-zero effective 1-cycles C_1 and C_2 such that

$$[C] = [C_1] + [C_2].$$

4.6. Proposition. Under the Assumption 4.1, the divisor $K_X + \Delta$ has the bend-and-break property.

The proof of this result needs some preparation:

4.7. Lemma. Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with rational singularities, and let Δ be a boundary divisor on X . Let $\mu : Y \rightarrow X$ be a bimeromorphic morphism from a normal $\mathbb{Q}$ -factorial threefold Y such that there exists a boundary divisor Δ_Y satisfying

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^*(K_X + \Delta).$$

Suppose that $K_Y + \Delta_Y$ has the bend-and-break property for some integer $d_{K_Y + \Delta_Y}$. Then $K_X + \Delta$ has the bend-and-break property.

Proof. Since $\dim X = 3$, there are at most finitely many curve in the image of the μ -exceptional locus E . Hence

$$d := \max\{d_{K_Y + \Delta_Y}, -(K_X + \Delta) \cdot Z \mid Z \text{ an irreducible curve s.t. } Z \subset \mu(E)\}$$

is a positive rational number. Let now $C \subset X$ be a curve such that

$$-(K_X + \Delta) \cdot C > d,$$

in particular $C \not\subset \mu(E)$. Thus the strict transform $\hat{C} \subset Y$ is well-defined and

$$-(K_Y + \Delta_Y) \cdot \hat{C} = -(K_X + \Delta) \cdot C > d \geq d_{K_Y + \Delta_Y}.$$

Consequently, there exist effective non-zero 1-cycles $\hat{C}_1$ and $\hat{C}_2$ on Y such that

$$[\hat{C}] = [\hat{C}_1] + [\hat{C}_2].$$

We claim that we can find a decomposition such that $\mu_*[\hat{C}_1]$ and $\mu_*[\hat{C}_2]$ are both non-zero. Since

$$[C] = \mu_*[\hat{C}] = \mu_*[\hat{C}_1] + \mu_*[\hat{C}_2]$$

this will finish the proof.

To prove the claim, fix a Kähler form ω_Y on Y such that $\omega_Y \cdot B \geq 1$ for every curve $B \subset Y$. We will prove the claim by induction on the degree $l := \omega_Y \cdot \hat{C}$. The start of the induction for $l = 1$ is trivial, since the class of a curve $\hat{C}$ with $\omega_Y \cdot \hat{C} = 1$ does not decompose. For the induction step suppose that the claim holds for every curve $B \subset Y$ with $\omega_Y \cdot B < l$ and $-(K_Y + \Delta_Y) \cdot B > d$. Suppose that $l \leq \omega_Y \cdot \hat{C} < l + 1$ and consider the decomposition

$$[\hat{C}] = [\hat{C}_1] + [\hat{C}_2].$$

If both $\mu_*[\hat{C}_1]$ and $\mu_*[\hat{C}_2]$ are non-zero, there is nothing to prove, so suppose that (up to renumbering) $\mu_*[\hat{C}_2] = 0$. Since $\hat{C}_2$ is effective, all the irreducible components of $\hat{C}_2$ are contracted by μ . Now $K_Y + \Delta_Y$ is μ -numerically trivial, so we get

$$-(K_Y + \Delta_Y) \cdot \hat{C}_1 = -(K_Y + \Delta_Y) \cdot \hat{C} > d.$$

Moreover since $\omega_Y \cdot \hat{C}_2 \geq 1$, we have $\omega_Y \cdot \hat{C}_1 < l$. Thus the induction hypothesis applies to C_1 , and since $[C] = \mu_*[\hat{C}] = \mu_*[\hat{C}_1]$ this proves the claim. $\square$

4.8. Lemma. *In the situation of Assumption 4.1, suppose that K_X has the bend-and-break property for some integer d_0 . Then $K_X + \Delta$ has the bend-and-break property.*

Proof. Since X is a threefold, the set $(\text{Supp } \Delta)_{\text{sing}} \cup X_{\text{sing}}$ is a finite union of curves and points. Hence

$$d := \max\{3, d_0, -(K_X + \Delta) \cdot Z \mid Z \text{ a curve s.t. } Z \subset (\text{Supp } \Delta)_{\text{sing}} \cup X_{\text{sing}}\}$$

is a positive rational number. Let now $C \subset X$ be a curve such that $-(K_X + \Delta) \cdot C > d$. If $\Delta \cdot C \geq 0$, then we have $-K_X \cdot C > d \geq d_0$, so we can apply the bend-and-break property for K_X . Therefore we may suppose

$$0 > \Delta \cdot C = \sum a_i \Delta_i \cdot C,$$

so, up to renumbering, we may suppose that $\Delta_1 \cdot C < 0$. Since $0 \leq a_i \leq 1$ (Δ is a boundary), this implies

$$-(K_X + \Delta_1 + \sum_{i \geq 2} a_i \Delta_i) \cdot C \geq -(K_X + \Delta) \cdot C > d.$$

Set $S := \Delta_1$ and $B := \sum_{i \geq 2} a_i \Delta_i$, and denote by $\pi : \hat{S} \rightarrow S$ the composition of the normalisation and the minimal resolution. Note that by definition of d , the curve C is not contained in the set $S_{\text{sing}} \cup (\sum_{i \geq 2} \Delta_i \cap S) \cup (X_{\text{sing}} \cap S)$. Therefore the strict transform $\hat{C} \subset \hat{S}$ is well-defined and (3) yields

$$-K_{\hat{S}} \cdot \hat{C} \geq -(K_X + \Delta_1 + \sum_{i \geq 2} a_i \Delta_i) \cdot C > d.$$

Since $d \geq 3$, an application of [HP13a, Lemma 5.5.b)] yields an effective 1-cycle $\sum_{k=1}^m \hat{C}_k$ in $\hat{S}$ with $m \geq 2$ such that

$$[\sum_{k=1}^m \hat{C}_k] = [\hat{C}],$$

such that $K_{\hat{S}} \cdot \hat{C}_1 < 0$, and $K_{\hat{S}} \cdot \hat{C}_2 < 0$. Since $K_{\hat{S}}$ is π_1 -nef, we conclude $\pi_*[\hat{C}_1] \neq 0$ and $\pi_*[\hat{C}_2] \neq 0$. Thus

$$[C] = \pi_*[\hat{C}] = \sum_{k=1}^m \pi_*[\hat{C}_k]$$

and the first two terms of this sum are non-zero. $\square$

Proof of Proposition 4.6. We will first prove the statement under some additional assumptions, then reduce the general case to this situation.

Step 1. Suppose that (X, Δ) is a lc pair and X has terminal singularities. By [HP13a, Cor.5.7] we know that K_X has the bend-and-break property. Thus by Lemma 4.8 the divisor $K_X + \Delta$ has the bend-and-break property.

Step 2. Suppose that X has klt singularities, i.e., $(X, 0)$ is klt. By Theorem 2.3 there exists a bimeromorphic morphism $\mu : Y \rightarrow X$ from a normal complex Kähler space Y with terminal singularities and a boundary divisor Δ_Y such that (Y, Δ_Y) is lc and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X.$$

By Step 1 and Lemma 4.7 the divisor K_X has the bend-and-break property.

Step 3. Suppose that (X, Δ) is dlt. Then $(X, 0)$ is klt [KM98, 2.39, 2.41]. By Step 2 and Lemma 4.8 this implies that $K_X + \Delta$ has the bend-and-break property.

Step 4. General case. By Theorem 2.4 there exists a bimeromorphic morphism $\mu : Y \rightarrow X$ and a boundary divisor Δ_Y on Y such that (Y, Δ_Y) is dlt and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^*(K_X + \Delta).$$

By Step 3 and Lemma 4.7 this implies that $K_X + \Delta$ has the bend-and-break property. $\square$

We can now prove the weak cone theorem for the classical Mori cone.

4.9. Proposition. *Under the Assumption 4.1 there exists a number $d \in \mathbb{Q}^+$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that*

$$0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$$

and

$$\overline{NE}(X) = \overline{NE}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i]$$

Proof. By Proposition 4.6 there exists a positive number $d \in \mathbb{Q}^+$ realising the bend-and-break property (cf. Definition 4.5) for $K_X + \Delta$. Since there are only countably many curve classes $[C] \in \overline{NE}(X)$, we may choose a representative Γ_i for each class such that $0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$. We set

$$V := \overline{NE}(X)_{(K_X + \Delta) \geq 0} + \sum_{0 < -(K_X + \Delta) \cdot \Gamma_i \leq d} \mathbb{R}^+[\Gamma_i].$$

Fix a Kähler form η_X on X such that $\eta_X \cdot C \geq 1$ for every curve $C \subset X$.

We need to prove that $\overline{NE}(X) = V$. By [HP13a, Lemma 6.1]⁴ it is sufficient to show that

$$\overline{NE}(X) = \overline{V},$$

i.e. the class $[C]$ of every irreducible curve $C \subset X$ is contained in V . We will prove the statement by induction on the degree $l := \eta_X \cdot C$. The start of the induction for $l = 0$ being trivial, we suppose that we have shown the statement for all curves of degree at most $l - 1$ and let C be a curve such that

$$l - 1 < \eta_X \cdot C \leq l.$$

⁴The statement in [HP13a, Lemma 6.1] is for the canonical class K_X , but the proof does not use this hypothesis.

If $-(K_X + \Delta) \cdot C \leq d$ we are done. Otherwise, there exists by the bend-and-break property a decomposition

$$[C] = [C_1] + [C_2]$$

with C_1 and C_2 non-zero effective 1-cycles on X . Since $\eta_X \cdot C_i \geq 1$ for $i = 1, 2$ we have $\eta_X \cdot C_i \leq l - 1$ for $i = 1, 2$. By induction both classes are in V , so $[C]$ is in V . $\square$

Proof of Theorem 4.2. We follow the strategy of the proof of [HP13a, Prop.6.4]. By Proposition 4.9 there exists a number $d \in \mathbb{Q}^+$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that $0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$ and

$$\overline{\text{NE}}(X) = \overline{\text{NE}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+ [\Gamma_i].$$

Set

$$V := \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+ [\Gamma_i].$$

By [HP13a, Lemma 6.1] it is sufficient to show that $\overline{\text{NA}}(X) \subset \overline{V}$. Let $\pi : \hat{X} \rightarrow X$ be a desingularisation, then by [HP13a, Prop.3.10] we have $\overline{\text{NA}}(X) = \pi_*(\overline{\text{NA}}(\hat{X}))$. Thus it is sufficient to prove that for a set of generators $\hat{\alpha}_i$ of $\overline{\text{NA}}(\hat{X})$, we have $\pi_*(\hat{\alpha}_i) \in \overline{V}$. By [DP04, Cor.0.3] the cone $\overline{\text{NA}}(\hat{X})$ is the closure of the convex cone generated by cohomology classes of the form $[\hat{\omega}]^2, [\hat{\omega}] \cdot [\hat{S}]$ and $[\hat{C}]$ where $\hat{\omega}$ is a Kähler form, $\hat{S}$ a surface and $\hat{C}$ a curve on $\hat{X}$. Let now $\hat{\alpha}$ be such a generator, then our goal is to show that the $\alpha := \pi_*(\hat{\alpha})$ of any of this three types is contained in $\overline{V}$.

1st case. $\hat{\alpha} = [\hat{\omega}]^2$ with $\hat{\omega}$ a Kähler form. Since $\pi^*(K_X + \Delta)$ is pseudoeffective, we have $\pi^*(K_X + \Delta) \cdot [\hat{\omega}]^2 \geq 0$, hence

$$(K_X + \Delta) \cdot \alpha = (K_X + \Delta) \cdot \pi_*(\hat{\alpha}) = \pi^*(K_X + \Delta) \cdot \hat{\alpha} \geq 0,$$

and thus $\alpha \in \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0}$.

2nd case. $\hat{\alpha} = [\hat{C}]$ with $\hat{C}$ a curve. Then set $C := \pi_*(\hat{C})$, so that $\alpha = [C]$. Since we have an inclusion

$$(13) \quad \overline{\text{NE}}(X) = \overline{\text{NE}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+ [\Gamma_i] \hookrightarrow \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+ [\Gamma_i],$$

and by hypothesis any curve class $[C]$ is in the left hand side, we see that $[C] \in V$.

(3) *3rd case.* $\hat{\alpha} = [\hat{\omega}] \cdot [\hat{S}]$ with $\hat{S}$ an irreducible surface and $\hat{\omega}$ a Kähler form. If

$$\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] \geq 0,$$

the class α is in $\overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0}$. Suppose now that

$$\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] < 0.$$

Using the projection formula we see that $\pi(\hat{S})$ is not a point.

Case a) Suppose that $\pi(\hat{S})$ is a surface S . Since $\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] < 0$, the restriction $\pi^*(K_X + \Delta)|_{\hat{S}}$ is not pseudoeffective. Thus the restriction $(K_X + \Delta)|_S$ is not pseudoeffective. Hence S is covered by rational curves by Lemma 4.3. Since $\hat{S} \rightarrow S$ is bimeromorphic, the same property holds for $\hat{S}$.

Let $\pi : \bar{S} \rightarrow \hat{S}$ be the composition of the normalisation and the minimal resolution, then $\bar{S}$ is a uniruled projective surface, in particular, $H^2(\bar{S}, \mathcal{O}_{\bar{S}}) = 0$. Hence the Chern class map

$$\text{Pic } \bar{S} \rightarrow H^2(\bar{S}, \mathbb{Z})$$

is surjective, so $\pi^*(\hat{\omega}|_{\hat{S}})$, which is a real closed form of type $(1, 1)$, is represented by a $\mathbb{R}$ -divisor which is nef and big. As in the proof of [HP13a, Prop.6.4], this implies that

$$\hat{\alpha} = [\hat{\omega}] \cdot [\hat{S}] \in \overline{\text{NE}}(\hat{X}).$$

Thus we have

$$\alpha = \pi_*([\hat{\omega}] \cdot [\hat{S}]) \in \overline{\text{NE}}(X),$$

so α is in the image of the inclusion (13).

Case b) Suppose that $\pi(\hat{S})$ is a curve C . We claim that there exists a number $\lambda \in \mathbb{R}$ such that

$$\alpha = \pi_*(\hat{\alpha}) = \lambda[C]$$

in $N_1(X)$, in particular α is in the image of the inclusion (13) which finishes the proof.

By duality it is sufficient to prove that there exists a $\lambda \in \mathbb{R}$ such that for every class $H \in N^1(X)$ we have

$$H \cdot \pi_*(\hat{\alpha}) = \lambda(H \cdot C).$$

By the projection formula

$$H \cdot \pi_*(\hat{\alpha}) = \pi^*H \cdot \hat{\alpha} = \pi^*H \cdot [\hat{\omega}] \cdot [\hat{S}] = [(\pi^*H)|_S] \cdot [\hat{\omega}|_{\hat{S}}].$$

By definition of C we have a surjective map $\hat{S} \rightarrow C$, so $[(\pi^*H)|_S]$ is numerically equivalent to $(H \cdot C)[F]$ where F is a general fibre of $\hat{S} \rightarrow C$. Thus we see that

$$[(\pi^*H)|_S] \cdot [\hat{\omega}|_{\hat{S}}] = \lambda(H \cdot C)$$

where $\lambda := [\hat{\omega}|_{\hat{S}}] \cdot F$ does not depend on H . $\square$

4.B. The contraction theorem. Suppose that the Assumption 4.1 holds. For the whole subsection we fix $R := \mathbb{R}^+[\Gamma_{i_0}]$ a $(K_X + \Delta)$ -negative extremal ray in $\overline{\text{NA}}(X)$. The following proposition is a well-known consequence of the weak cone theorem 4.2, cf. [HP13a, Prop.7.3] for details:

4.10. Proposition. *There exists a nef class $\alpha \in N^1(X)$ such that*

$$R = \{z \in \overline{\text{NA}}(X) \mid \alpha \cdot z = 0\},$$

and such that, using the notation of Theorem 4.2, the class α is strictly positive on

$$\left(\overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I, i \neq i_0} \mathbb{R}^+[\Gamma_i] \right) \setminus \{0\}.$$

We call α a nef supporting class for the extremal ray R .

Note first that by hypothesis, the cohomology class $\alpha - (K_X + \Delta)$ is positive on the extremal ray R , moreover we know that α is positive on

$$\left(\overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I, i \neq i_0} \mathbb{R}^+[\Gamma_i] \right) \setminus \{0\}.$$

Thus, up to replacing α by some positive multiple, we can suppose that $\alpha - (K_X + \Delta)$ is positive on $\overline{NA}(X) \setminus \{0\}$. Since X is a Kähler space, this implies by [HP13a, Cor.3.14] that

$$(14) \quad \omega := \alpha - (K_X + \Delta)$$

is a Kähler class. Thus the nef class $\alpha = (K_X + \Delta) + \omega$ is big, the divisor $K_X + \Delta$ being $\mathbb{Q}$ -effective. In particular we have

$$(15) \quad \alpha^3 > 0.$$

We divide extremal rays into two classes, according to the deformation behaviour of the curves they contain:

4.11. Definition. *We say that the $(K_X + \Delta)$ -negative extremal ray R is small if every curve $C \subset X$ with $[C] \in R$ is very rigid in the sense of [HP13a, Defn.4.3]. Otherwise we say that the extremal ray R is divisorial.*

4.B.1. *Small rays.*

4.12. Theorem. *Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_{i_0}]$ is small. Then the contraction of the ray R exists.*

The proof of this theorem will be a refinement of the argument in [HP13a] combined with a recent result by Collins and Tosatti [CT13]. The following lemma is the key ingredient:

4.13. Proposition. *Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_i]$ is small. Let $S \subset X$ be an irreducible surface. Then we have $\alpha^2 \cdot S > 0$.*

At this point we cannot yet exclude the possibility that there are infinitely many distinct curves $C \subset X$ such that $[C] \in R$ ⁵. However by definition of a small ray no such curve (or its multiples) deforms. Since the irreducible components of the cycle space are countable we see that there are at most countably many curves $C \subset X$ such that $[C] \in R$.

Proof of Proposition 4.13. If $\alpha|_S = 0$ then (14) implies that

$$-(K_X + \Delta)|_S = \omega|_S.$$

Thus the divisor $-(K_X + \Delta)|_S$ is ample, in particular S is projective and S is covered by curves. Since $\alpha|_S = 0$, the classes of all these curves are in R , a contradiction.

Arguing by contradiction we suppose now that $\alpha|_S \neq 0$ but $\alpha^2 \cdot S = 0$. Then we have

$$0 = \alpha^2 \cdot S = (K_X + \Delta) \cdot \alpha \cdot S + \omega \cdot \alpha \cdot S$$

and

$$\omega \cdot \alpha \cdot S = \omega|_S \cdot \alpha|_S > 0$$

by the Hodge index theorem. Thus we obtain

$$(16) \quad (K_X + \Delta) \cdot \alpha \cdot S < 0.$$

In particular $(K_X + \Delta)|_S$ is not pseudoeffective, the class $\alpha|_S$ being nef.

⁵If X has terminal singularities we can use some additional argument to obtain this property, cf. [HP13a, Rem.7.2].

Let $\pi : \hat{S} \rightarrow S$ be the composition of normalisation and minimal resolution (cf. Subsection 2.B). We claim that there exists an effective $\mathbb{Q}$ -divisor E on $\hat{S}$ such that

$$(17) \quad (K_{\hat{S}} + E) \cdot \pi^*(\alpha|_S) < 0.$$

Assuming this for the time being, let us see how to derive a contradiction: note first that $K_{\hat{S}}$ is not pseudoeffective, so $\hat{S}$ is uniruled and projective. In particular the nef class $\pi^*(\alpha|_S)$ is represented by an $\mathbb{R}$ -divisor. Fix an ample divisor A on $\hat{S}$. By [Ara10, Thm.1.3] we know that given $\varepsilon > 0$, there exists a decomposition

$$\pi^*(\alpha|_S) = C_\varepsilon + \sum \lambda_{i,\varepsilon} M_{i,\varepsilon}$$

where $\lambda_{i,\varepsilon} \geq 0$, $(K_{\hat{S}} + \varepsilon A) \cdot C_\varepsilon \geq 0$ and the $M_{i,\varepsilon}$ are movable curves. Since $M_{i,\varepsilon}$ belongs to an (uncountable) deformation family of curves we obtain

$$\pi^*(\alpha|_S) \cdot M_{i,\varepsilon} = \alpha \cdot \pi_*(M_{i,\varepsilon}) > 0.$$

Since $(\pi^*(\alpha|_S))^2 = 0$ this implies that $\pi^*(\alpha|_S) = C_\varepsilon$ for all $\varepsilon > 0$. Passing to the limit we obtain $K_{\hat{S}} \cdot \pi^*(\alpha|_S) \geq 0$, a contradiction to (17).

Proof of the claim (17). As in the proof of Lemma 4.3 we need a case distinction.

1st case. Suppose that $K_X \cdot \alpha \cdot S < 0$. Since K_X is $\mathbb{Q}$ -effective, S is one of the surfaces S_j in the decomposition (12) and $S \cdot \alpha \cdot S < 0$. In particular we have

$$(18) \quad (K_X + S) \cdot \alpha \cdot S < 0.$$

Applying (2) with $B = 0$ we obtain an effective $\mathbb{Q}$ -divisor E on $\hat{S}$ such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S)|_S$. Thus (17) follows from (18).

2nd case. Suppose that $K_X \cdot \alpha \cdot S \geq 0$. By (16) this implies that $\Delta \cdot \alpha \cdot S < 0$, so S is contained in the support of Δ ; up to renumbering we may suppose that $S = \Delta_1$. Since $\Delta_i \cdot \alpha \cdot S \geq 0$ for every $i \geq 2$, it follows $S \cdot \alpha \cdot S < 0$. Since $0 \leq a_i \leq 1$ we conclude

$$(19) \quad (K_X + S + \sum_{i \geq 2} a_i \Delta_i) \cdot \alpha \cdot S \leq (K_X + \Delta) \cdot \alpha \cdot S < 0.$$

Applying (2) with $B = \sum_{i \geq 2} a_i \Delta_i$ we obtain an effective $\mathbb{Q}$ -divisor E on $\hat{S}$ such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S + \sum_{i \geq 2} a_i \Delta_i)|_S$. Thus (17) follows from (19). $\square$

Proof of Theorem 4.12. Let $\pi : \hat{X} \rightarrow X$ be a desingularisation. Since α is nef and big, the pull-back $\pi^*\alpha$ is nef and big. By [CT13, Thm.1.1] the non-Kähler locus $E_{nK}(\pi^*\alpha)$ is equal to the null-locus, i.e. we have

$$E_{nK}(\pi^*\alpha) = \cup_{(\pi^*\alpha)|_Z \equiv 0} Z,$$

where the union runs over all the subvarieties of $\hat{X}$. If $Z \subset X$ is a surface such that $(\pi^*\alpha)|_Z^2 = 0$ then it follows from the projection formula and Proposition 4.13 that $\dim \pi(Z) \leq 1$. Thus we see that $E_{nK}(\pi^*\alpha)$ is a finite union of π -exceptional surfaces and curves. Since

$$E_{nK}(\alpha) \subset \pi(E_{nK}(\pi^*\alpha)),$$

and X_{sing} being a union of curves and points, we see that $E_{nK}(\alpha)$ is a union of curves and points⁶. Clearly $E_{nK}(\alpha)$ contains all the curves $C \subset X$ such that

⁶If X has isolated singularities, the argument shows that

$$E_{nK}(\alpha) = \cup_{\alpha \cdot B = 0} B.$$

$[C] \in R$, but it is not clear that equality holds. However the arguments of the proof of [HP13a, Thm.7.14] work exactly in this situation, so we can contract the curves in the extremal ray. $\square$

4.B.2. Divisorial rays.

4.14. Notation. Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_{i_0}]$ is divisorial. Since the divisor $K_X + \Delta$ is $\mathbb{Q}$ -effective and $(K_X + \Delta) \cdot R < 0$ there exists an irreducible surface $S \subset X$ such that $S \cdot R < 0$. In particular any curve $C \subset X$ with $[C] \in R$ is contained in S and S is covered by these curves.

Let $\nu : \tilde{S} \rightarrow S \subset X$ be the normalisation; then $\nu^*(\alpha|_S)$ is a nef class on $\tilde{S}$ and we may consider the nef reduction

$$\tilde{f} : \tilde{S} \rightarrow \tilde{T}$$

with respect to $\nu^*(\alpha|_S)$, cf. [BCE⁺02, Thm.2.6]. Since S is covered by curves that are α -trivial, the surface $\tilde{S}$ is covered by curves that are $\nu^*(\alpha|_S)$ -trivial. By definition of the nef reduction this implies

$$n(\alpha) := n(\nu^*(\alpha|_S)) := \dim \tilde{T} \in \{0, 1\}.$$

4.15. Lemma. *Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 0$. Then the surface S can be blown down to a point p : there exists a birational morphism $\varphi : X \rightarrow Y$ to a normal compact threefold Y with $\dim \varphi(S) = 0$ such that $\varphi|_{X \setminus S}$ is an isomorphism onto $Y \setminus \{p\}$.*

Proof. The proof is identical to the proof of [HP13a, Cor.7.7] which only uses that $S \cdot R < 0$. $\square$

The case $n(\alpha) = 1$ is much more subtle. If $S \subset \text{Supp } \Delta$, we can suppose up to renumbering that $S = \Delta_1$ and set $B = \sum_{i \geq 2} a_i \Delta_i$. If $S \not\subset \text{Supp } \Delta$ we simply set $B = \Delta$. In both cases we have by (1) that

$$(20) \quad K_{\tilde{S}} + B_{\tilde{S}} \sim_{\mathbb{Q}} \nu^*(K_X + S + B)|_S,$$

where $B_{\tilde{S}}$ is a canonically defined effective $\mathbb{Q}$ -divisor. Let $\tilde{C}$ be a general fibre of the nef reduction $\tilde{f} : \tilde{S} \rightarrow \tilde{T}$. Since $[\nu(\tilde{C})] \in R$ we have

$$(K_X + \Delta) \cdot \nu(\tilde{C}) < 0 \quad \text{and} \quad S \cdot \nu(\tilde{C}) < 0.$$

Since $0 \leq a_1 \leq 1$ this implies

$$(K_X + S + B) \cdot \nu(\tilde{C}) \leq (K_X + \Delta) \cdot \nu(\tilde{C}) < 0.$$

Thus we have $(K_{\tilde{S}} + B_{\tilde{S}}) \cdot \tilde{C} < 0$, in particular $\tilde{C}^2 = 0$ implies that $\tilde{C} \simeq \mathbb{P}^1$. The normal surface $\tilde{S}$ being smooth in a neighbourhood of the fibre $\tilde{C}$, we conclude $K_{\tilde{S}} \cdot \tilde{C} = -2$. Since $B_{\tilde{S}} \cdot \tilde{C} \geq 0$ we arrive at

4.16. Lemma. *Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 1$. Then the extremal ray R is represented by a rational curve C such that*

$$-(K_X + \Delta) \cdot C \leq 2.$$

This estimate allows us to complete the proof of the cone theorem:

This allows to simplify the proof in [HP13a].

Proof of Theorem 1.3. The only statement that is not part of Theorem 4.2 is that in every $(K_X + \Delta)$ -negative extremal ray R_i we can find a rational curve Γ_i , and, in case (X, Δ) is dlt, such that $\Gamma_i \in R_i$ and

$$0 < -(K_X + \Delta) \cdot \Gamma_i \leq 4.$$

If the extremal ray Γ_i is divisorial with $n(\alpha) = 1$, we conclude by Lemma 4.16, even if (X, Δ) is only lc. If the extremal ray Γ_i is small or divisorial with $n(\alpha) = 0$, the contraction exists by Theorem 4.12 and Lemma 4.15. Since (X, Δ) is dlt, we may simply apply [Kaw91] [Deb01, Thm.7.46]. $\square$

4.17. Remark. In order to prove Theorem 1.3 in full generality in the lc case, it remains to prove the existence of rational curves if the ray is divisorial contracting a divisor S to a point or if the ray is small, with the bound $d = 4$. The existence of rational curves in the first case just follows from the arguments preceding Lemma 4.16, which show that S is uniruled. The existence of rational curves in the small case requires some vanishing theorem which is not yet established in the Kähler case: if W is the union of the lc centers, then $H^1(W, \mathcal{O}_W) = 0$.

In order to prove the existence of the contraction in the case $n(\alpha) = 1$ we would like to construct a fibration $S \rightarrow T$ whose normalisation is the nef reduction $\tilde{S} \rightarrow \tilde{T}$. At the moment we can only realise this strategy under an additional condition on the singularities of S . We will use the notion of a semi-log-canonical (slc) surface; see sect.5.

4.18. Proposition. *Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 1$. Suppose moreover S has slc singularities, i.e. $(S, 0)$ is slc. Then there exists a morphism with connected fibres $S \rightarrow T$ onto a curve T that contracts a curve $C \subset S$ if and only if $\alpha|_S \cdot C = 0$. Moreover the contraction of the extremal ray R exists.*

Proof. Since $(S, 0)$ is slc, it has normal crossing singularities in codimension one. The normalisation map $\nu : \tilde{S} \rightarrow S$ is finite, so the general curve $\nu(\tilde{C})$ is contained in the locus where S has normal crossing singularities. If $\nu(\tilde{C})$ is in $S_{\text{non-s}}$ the proof of [HP13a, Lemma 7.8] applies without changes, so suppose that this is not the case. Let

$$Z = \cup Z_k \subset S_{\text{sing}}$$

be the union of curves Z_k such that $\nu^{-1}(Z_k)$ meets the general curve $\tilde{C}$. For each k we set $\tilde{Z}_k := \nu^{-1}(Z_k)$ and set $\tilde{Z} := \cup \tilde{Z}_k$. Since S has normal crossings in the generic point of Z_k , the natural map $\tilde{Z}_k \rightarrow Z_k$ has degree two. Note now that every irreducible component of $\tilde{Z}$ is $\nu^*\alpha$ -positive: otherwise $\tilde{S}$ would be connected by $\nu^*\alpha$ -trivial curves, thus $n(\alpha) = 0$ in contradiction to our assumption. In particular every curve in $\tilde{Z}$ dominates $\tilde{T}$.

Since $\tilde{Z}$ maps into the normal crossings locus of S we can decompose the differend (20)

$$B_{\tilde{S}} = \tilde{Z} + R$$

where R is an effective $\mathbb{Q}$ -divisor with no common component with $\tilde{Z}$. The intersection points of $\tilde{Z}$ and $\tilde{C}$ are contained in the smooth locus of $\tilde{S}$, so $\tilde{Z} \cdot \tilde{C}$ is a positive integer. On the other hand we know that $K_{\tilde{S}} \cdot \tilde{C} = -2$ and

$$-2 < (K_{\tilde{S}} + \tilde{Z}) \cdot \tilde{C} \leq (K_{\tilde{S}} + \tilde{Z} + R) \cdot \tilde{C} < 0.$$

Thus $\tilde{Z} \cdot \tilde{C} = 1$. Since all the irreducible components of $\tilde{Z}$ surject onto $\tilde{T}$, this shows that $\tilde{Z}$ is irreducible. In particular Z itself is irreducible. Moreover we conclude that

- a) the curves Z and C meet in a unique point q ; and
- b) we have $\nu^{-1}(q) = \{p_1, p_2\}$ with $p_1 \in \tilde{C}$, but $p_2 \notin \tilde{C}$.

Since $\tilde{C}$ is general and ν is finite, the point p_2 lies on another general fibre $\tilde{C}$ such that $\nu(\tilde{C})$ and $\nu(\tilde{C})$ meet in q . Let now T be the unique irreducible component of the cycle space $\text{Chow}(S)$ such that the general point corresponds to the cycle $\nu(\tilde{C}) + \nu(\tilde{C})$, and let $\Gamma \rightarrow T'$ be the semi-normalisation [Kol07, p.156] of the universal family over T . By construction we have a natural bimeromorphic morphism $\Gamma \rightarrow S$ and, by what precedes, this morphism is an isomorphism in the neighbourhood of the general fibre F of $\Gamma \rightarrow T'$. Thus F defines a Cartier divisor on S such that $F^2 = 0$ and $\kappa(S, F) = 1$. Thus some positive multiple of F defines a morphism with connected fibres $S \rightarrow T$ and one easily checks that this morphism realises the nef reduction with respect to $\alpha|_S$. Now conclude as in [HP13a, Cor.7.9]. $\square$

4.C. Running the MMP. We first collect a number of results which we could not find in this form in the literature and give an indication how to adapt the proof.

4.19. Theorem. ([Sho92], in the algebraic case [Kwc92]) *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold. Let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray R in $\overline{NA}(X)$. Suppose that the contraction of $\varphi : X \rightarrow Y$ is small. Then the flip $\varphi^+ : X^+ \rightarrow Y$ exists. Moreover X^+ is a normal $\mathbb{Q}$ -factorial compact Kähler threefold and $(X^+, (\varphi^+)^{-1}_* \circ \varphi_* \Delta)$ is dlt.*

Proof. Since X is $\mathbb{Q}$ -factorial, the pair $(X, (1 - \varepsilon)\Delta)$ is klt for every $0 < \varepsilon < 1$. Moreover for $0 < \varepsilon \ll 1$ the divisor $K_X + (1 - \varepsilon)\Delta$ is negative on the extremal ray R . Thus the flip $\varphi : X^+ \rightarrow Y$ exists by [Sho92, Thm], see also [Cor07] (and [Kwc92, 2.32] for passing to the limit). We alternatively may apply [Kwc92], where the existence of log flips is reduced to the existence of terminal flips, [Mor88]. The existence of terminal flips is now a local analytic construction. The reduction to the terminal case as given in sections 5,6 and 8 in [Kwc92] works in the analytic setting as well⁷. By Corollary 3.9, the complex space Y is Kähler. Since the morphism φ^+ is projective, X^+ is again Kähler. For the remaining properties we refer to Proposition 3.6. $\square$

While the existence of flips is a local property [KM98, 6.7], this is not necessarily the case for termination results. However in most cases we will only use special termination for dlt pairs which is much simpler to prove:

4.20. Theorem. [Sho92, Kwc92] *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold. Let Δ be an effective reduced Weil divisor such that the pair (X, Δ) is dlt. Set $(X_0, \Delta_0) := (X, \Delta)$ and let*

$$(\varphi_i : (X_i, \Delta_i) \dashrightarrow (X_{i+1}, \Delta_{i+1} := (\varphi_i)_* \Delta_i))_{i \in I}$$

be a sequence of $(K + \Delta)$ -flips where $I \subset \mathbb{N}$. Then for all $i \gg 0$ the flipping locus is disjoint from Δ_i .

⁷It is important to notice that the construction is locally analytic near the contracted curves.

In particular, if for every $i \in I$ the flipping locus of φ_i is contained in Δ_i , then the sequence of $(K + \Delta)$ -flips terminates.

Proof. The proof of [Fuj07, Thm.4.2.1] applies without changes: obviously the MMP exists for normal $\mathbb{Q}$ -factorial compact Kähler surfaces, the discrepancy calculations are local properties, so they also apply in the non-algebraic setting. $\square$

4.21. Theorem. [Kaw92b, Kwc92] *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold such that $(X, 0)$ is klt. Set $X_0 := X$ and let*

$$(\varphi_i : X_i \dashrightarrow X_{i+1})_{i \in I}$$

be a sequence of K -flips. Then I is finite, that is any sequence of K -flips terminates.

Proof. The proof of [Kaw92b, Thm.1] is based on three tools:

- a) existence of terminal models for klt pairs [Kaw92b, Thm.5];
- b) crepant extraction of a unique divisor with nonpositive discrepancy [Kaw92b, Lemma 6];
- c) discrepancy calculations.

The first two tools are available in the Kähler setting since we can argue as in Subsection 2.A. Discrepancy calculations are local properties, so they also apply in the non-algebraic setting. $\square$

As an application we can run the MMP for certain dlt pairs:

4.22. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold that is not uniruled. Let $D \in |mK_X|$ be a pluricanonical divisor and set $B := \text{Supp } D$. Suppose that the pair (X, B) is dlt. Then there exists a terminating $(K + B)$ -MMP, that is, there exists a birational map*

$$\varphi : (X, B) \dashrightarrow (X', B' := \varphi_* B)$$

which is a composition of $K + B$ -negative divisorial contractions and flips such that X' is a normal $\mathbb{Q}$ -factorial compact Kähler threefold, the pair (X', B') is dlt and $K_{X'} + B'$ is nef.

Proof. Set $(X_0, B_0) := (X, B)$ and $D_0 := D$.

Step 1. Existence of the MMP. If $K_{X_i} + B_i$ is nef, there is nothing to prove, so suppose that this is not the case. Then there exists by Theorem 1.3 a $K_{X_i} + B_i$ -negative extremal ray $R_i \in \overline{\text{NA}}(X_i)$. We may apply Theorem 1.3, since dlt pairs have rational singularities, [KM98, 5.22]. Since X_i is not uniruled, the extremal ray is divisorial or small.

If the extremal ray is small, the contraction $\psi_i : X_i \rightarrow Y_i$ exists by Theorem 4.12. By Theorem 4.19 the flip $\psi_i^+ : X_i^+ \rightarrow Y_i$ exists, and we denote by

$$\varphi_i : (X_i, B_i) \dashrightarrow (X_{i+1} := X_i^+, B_{i+1} := (\varphi_i)_* B_i)$$

the composition $(\psi_i^+)^{-1} \circ \psi_i$. By Theorem 4.19 we know that X_{i+1} is a normal $\mathbb{Q}$ -factorial Kähler space and (X_{i+1}, B_{i+1}) is dlt, so we can continue the MMP. Note finally that $D_{i+1} := (\varphi_i)_* D_i$ is a pluricanonical divisor such that $\text{Supp } D_{i+1} = B_{i+1}$. If the extremal ray is divisorial, let S_i be the unique surface such that $S_i \cdot R_i < 0$ (cf. Notation 4.14). Since $K_{X_i} + B_i$ is $\mathbb{Q}$ -effective and represented by an effective $\mathbb{Q}$ -divisor with support in B_i , the surface S_i must be an irreducible component of

B_i . Since the pair (X_i, B_i) is dlt, any of the irreducible components of B_i is normal [KM98, 5.52]. Moreover, there exists a boundary divisor $B_{i,S}$ such that

$$K_{S_i} + B_{i,S} \sim_{\mathbb{Q}} K_{X_i} + B_i$$

and the pair $(S_i, B_{i,S})$ is slc [Kwc92, 16.9.1]. Thus by Proposition 4.18 the contraction

$$\varphi_i : (X_i, B_i) \rightarrow (X_{i+1}, B_{i+1} := (\varphi_i)_* B_i)$$

exists and X_{i+1} is a Kähler space by Corollary 3.9. Moreover by Proposition 3.6, (X_{i+1}, B_{i+1}) is dlt and X_{i+1} is $\mathbb{Q}$ -factorial, so we can continue the MMP. Note finally that $D_{i+1} := (\varphi_i)_* D_i$ is a pluricanonical divisor such that $\text{Supp } D_{i+1} = B_{i+1}$.

Step 2. Termination. For every $i \in I$ we have $\text{Supp } D_i = B_i$, hence $K_{X_i} + B_i$ is $\mathbb{Q}$ -effective and represented by an effective $\mathbb{Q}$ -divisor with support in B_i . In particular the support of the extremal contraction is in B_i . Thus any sequence of $(K+B)$ -flips terminates by Theorem 4.20. $\square$

5. SEMI-LOG CANONICAL KÄHLER SURFACES

In this short section we gather the results concerning abundance in dimension 2 which will be used in the concluding sections. We refer to [Kwc92, Ch.12] [Kol13, Ch.5] for basic definitions of the theory of semi-log canonical (slc) surfaces. All the statements in this section are shown for projective surfaces in [Kwc92, Ch.12], we will see that the proofs also apply to our case with some minor modifications.

Let S be a reduced compact complex surface. We say that S is in class $\mathcal{C}$ if the desingularisations of the irreducible components S_i of S are Kähler. The Kähler property is a bimeromorphic invariant of smooth compact surfaces, so this definition is independent of the choice of the desingularisations.

5.1. Proposition. *Let S be an irreducible reduced compact complex surface in class $\mathcal{C}$. Let Δ be an effective $\mathbb{Q}$ -Cartier divisor on S such that the pair (S, Δ) is slc. Suppose that $K_S + \Delta$ is nef and numerically trivial. Then $K_S + \Delta$ is torsion, i.e. there exists a $m \in \mathbb{N}$ such that*

$$\mathcal{O}_S(m(K_S + \Delta)) \simeq \mathcal{O}_S.$$

Proof. Let $\nu : \tilde{S} \rightarrow S$ be the normalisation, and let $\tilde{\Delta}$ be the differend (cf. Subsection 2.B), so that we have

$$K_{\tilde{S}} + \tilde{\Delta} \sim_{\mathbb{Q}} \nu^*(K_S + \Delta) \equiv 0,$$

and the pair $(\tilde{S}, \tilde{\Delta})$ is lc.

1st case. $\tilde{\Delta} \neq 0$. In this case the surface $\tilde{S}$ is projective: indeed the anticanonical divisor $-K_{\tilde{S}} \equiv \tilde{\Delta}$ is pseudoeffective, so a K -MMP⁸ $\tilde{S} \rightarrow \bar{S}$ terminates with a Mori fibre space $\bar{S} \rightarrow W$. Since $\bar{S}$ is a surface and $-K_{\bar{S}}$ is relatively ample we see that $\bar{S}$ is projective. Since $\tilde{S} \rightarrow \bar{S}$ is a projective morphism, the surface $\tilde{S}$ is projective. Thus the statements in [Kwc92, Ch.12] apply.

2nd case. $\tilde{\Delta} = 0$. In this case S is normal, so if $\pi : \hat{S} \rightarrow S$ is the minimal resolution, then we have

$$\kappa(\hat{S}, K_{\hat{S}} + E) = \kappa(S, K_S),$$

⁸The existence of a MMP for a compact Kähler surface with lc singularities can be established following our arguments in Section 4.

where E is the canonical defined effective divisor such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* K_S \equiv 0$. Now if $E \neq 0$, then the smooth surface $\hat{S}$ is uniruled, hence projective, and therefore $\kappa(\hat{S}, K_{\hat{S}} + E) = 0$ by [Kwc92, Thm.12.1.1]. If $E = 0$, then S is a compact Kähler surface with numerically trivial canonical bundle, so $\kappa(\hat{S}, K_{\hat{S}}) = 0$ by the classification of compact surfaces [BHPVdV04]. $\square$

5.2. Lemma. *Let S be a normal compact Kähler surface, and let Δ be a boundary divisor on S . Suppose that the pair (S, Δ) is lc, that the class $K_S + \Delta$ is nef and that $\nu(K_S + \Delta) = 1$. Then $K_S + \Delta$ is semi-ample.*

Proof. Let $\mu : S' \rightarrow S$ be a dlt model, cp. Theorem 2.4, then (S', Δ') is dlt and $K_{S'} + \Delta' \sim_{\mathbb{Q}} \mu^*(K_S + \Delta)$. In particular $K_S + \Delta$ is semiample if and only if $K_{S'} + \Delta'$ is semiample. Thus we can assume without loss of generality that (S, Δ) is dlt.

Dlt singularities are rational [KM98, Thm.5.22], so the surface S is projective if and only if some desingularisation is projective [Nam02]. Thus if the algebraic dimension $a(S)$ is two, we conclude by [Kwc92, Thm.12.1.1]. If $a(S) = 1$, let $f : S \rightarrow C$ be the algebraic reduction. The general fibre F is an elliptic curve, so $K_S|_F \simeq \mathcal{O}_F$. Moreover every curve in S is contracted by f , so $\Delta \cap F = \emptyset$. Since $K_S + \Delta$ is nef, this implies

$$K_S + \Delta \sim_{\mathbb{Q}} f^* A,$$

with A an ample $\mathbb{Q}$ -divisor on C . Thus $K_S + \Delta$ is semiample. Finally we want to exclude the case $a(S) = 0$: note first that S contains only finitely many curves, so a desingularisation $\hat{S} \rightarrow S$ is not uniruled. Thus we have

$$\kappa(K_S) \geq \kappa(K_{\hat{S}}) \geq 0,$$

in particular $K_S + \Delta$ is $\mathbb{Q}$ -linearly equivalent to an effective divisor. Yet the intersection form on a surface with $a(S) = 0$ is negative definite, hence $(K_S + \Delta)^2 < 0$. In particular $K_S + \Delta$ is not nef, a contradiction. $\square$

5.3. Proposition. *Let S be a connected reduced compact complex surface in class $\mathcal{C}$. Let Δ be an effective $\mathbb{Q}$ -Cartier divisor on S such that the pair (S, Δ) is slc. Suppose that $K_S + \Delta$ is nef with numerical dimension $\nu(K_S + \Delta) \leq 1$. Suppose also that for every irreducible component $T \subset S$ the restriction $(K_S + \Delta)|_T$ is not zero.*

Then the divisor $K_S + \Delta$ is semi-ample.

Proof. Let $\nu : T \rightarrow S$ be the composition of a minimal semi-resolution (cf. [Kwc92, Defn.12.2.1, Prop.12.2.3]) with the normalisation. Since (S, Δ) is slc there exists for every irreducible component $T_i \subset T$ a boundary divisor Δ_i such that

$$K_{T_i} + \Delta_i \sim_{\mathbb{Q}} \nu^*(K_S + \Delta).$$

By our assumption $K_{T_i} + \Delta_i$ is nef of numerical dimension one, so $K_{T_i} + \Delta_i$ is semiample by Lemma 5.2. The whole point is now to prove that sufficiently many pluricanonical sections descend to S . However the proof of this descent theorem in [Kwc92, Ch.12.4] is based on [Kwc92, Prop.12.3.2, Thm.12.3.4] which are statements about proper morphisms. Thus they apply in our setting. $\square$

5.4. Theorem. *Let S be a compact connected reduced complex space of dimension two in class $\mathcal{C}$ with slc singularities. Then the natural maps induced by $\mathbb{C} \subset \mathcal{O}_S$*

$$H^p(S, \mathbb{C}) \rightarrow H^p(S, \mathcal{O}_S)$$

are surjective for every $p \in \mathbb{N}$.

Proof. Simply note that the proof of Theorem 12.1.2 of [Kwc92] works practically word by word. The main ingredient [Kwc92, 12.2.8] is about germs of slc surfaces and works in our situation. $\square$

6. ABUNDANCE: REDUCTION STEPS

This section corresponds to the preparatory lemmas in [Kwc92, Ch.13, Ch.14]. While the statements and the basic strategy of proof are quite similar we have to be more careful since we do not have a full log-MMP at our disposal. Moreover we make several points more precise which allows us to conclude quicker in Section 8.

6.1. Lemma. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with terminal singularities. Suppose that K_X is nef and $\nu(X) > 0$. Then there exists a normal $\mathbb{Q}$ -factorial compact Kähler threefold X' that is bimeromorphic to X and a $D' \in |mK_{X'}|$ with the following properties:*

- a) *Set $B' := \text{Supp } D'$. Then the pair (X', B') is dlt and $X' \setminus B'$ has terminal singularities.*
- b) *The divisor $K_{X'} + B'$ is nef and we have*
 - (i) $\nu(X) = \nu(K_{X'} + B')$; and
 - (ii) $\kappa(X) = \kappa(K_{X'} + B')$.

The proof needs some technical preparation:

6.2. Remark. Let X be a normal $\mathbb{Q}$ -factorial compact Kähler space, and let $D \in |mK_X|$ be an effective pluricanonical divisor. If we set $B = \text{Supp } D$, then

$$\kappa(K_X) = \kappa(K_X + B).$$

Indeed, B being effective, the inequality $\kappa(K_X) \leq \kappa(K_X + B)$ is obvious. Yet we also have an inclusion of effective divisors $B \subset D$, so we get

$$\kappa(K_X + B) \leq \kappa(K_X + D) = \kappa((m+1)K_X) = \kappa(X).$$

The following basic lemma has been shown in the algebraic case in [Kwc92, 13.2.4].

6.3. Lemma. *Let $\psi : V \rightarrow W$ be a bimeromorphic proper Kähler morphism between normal complex spaces. Let D be an effective $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor that is ψ -nef. Then we have*

$$\text{Supp } \psi_*(D) = \psi(\text{Supp } D),$$

i.e., the image of D has pure codimension one.

Proof. The inclusion $\text{Supp } \psi_*(D) \subset \psi(\text{Supp } D)$ is trivial. Fix now a point $w \in \psi(\text{Supp } D) \subset W$. Arguing by contradiction we assume that all the irreducible components of $\psi(\text{Supp } D)$ passing through w have dimension at most $\dim W - 2$. Thus, up to replacing W by an analytic neighbourhood of w , we assume that $\psi(\text{Supp } D)$ has dimension at most $\dim W - 2$. Then $\psi_*(-D) = 0$ and $-(-D)$ is ψ -nef, contradicting the negativity lemma, see e.g. [BCHM10, 3.6.2], unless $D = 0$. $\square$

The following crucial lemma seems to be well-known to experts and is used in several places without further mentioning, but we prefer to write it down in detail:

6.4. Lemma. *Let $X =: X_0$ be a normal $\mathbb{Q}$ -factorial compact Kähler space, and let $M =: M_0$ be an effective Cartier divisor on X that is nef. Let $N =: N_0$ be an effective Cartier divisor such that $\text{Supp } M = \text{Supp } N$, and let*

$$(\varphi_i : X_i \dashrightarrow X_{i+1})_{i=0, \dots, n}$$

be a finite sequence of N -negative contractions, that is every X_i is a normal $\mathbb{Q}$ -factorial compact Kähler space and φ_i is the divisorial contraction or flip of an extremal ray $R_i \in \overline{NA}(X_i)$ that is N_i -negative⁹. Then for every $i = 0, \dots, n$ the N -MMP induces an isomorphism

$$(X_0 \setminus \text{Supp } M_0) \simeq (X_{i+1} \setminus \text{Supp } M_{i+1}).$$

Remark. The essence of Lemma 6.4 is the following. Given the flip $\varphi : X_i \dashrightarrow X_{i+1}$ with contractions $f_i : X_i \rightarrow Y_i$ and $f_i^+ : X_{i+1} \rightarrow Y_i$, then the exceptional locus of f_i^+ is contained in M_{i+1} .

Proof. For every $i = 1, \dots, n$ we set $M_{i+1} := (\varphi_i)_* M_i$. Moreover we denote by Γ_i the normalisation of the graph of φ_i and by $p_i : \Gamma_i \rightarrow X_i$ and $q_i : \Gamma_i \rightarrow X_{i+1}$ the natural maps.

We will construct inductively a sequence of normal compact Kähler spaces V_i admitting bimeromorphic morphisms $f_i : V_i \rightarrow X$ and $g_i : V_i \rightarrow X_{i+1}$ such that

- a) the support of $f_i^* M$ contains the g_i -exceptional locus;
- b) g_i factors through Γ_i , that is there exists a bimeromorphic map $\beta_i : V_i \rightarrow \Gamma_i$ such that $g_i = q_i \circ \beta_i$.

Assuming this for the time being, let us see how to conclude: arguing by induction we suppose that we have an isomorphism

$$(X_0 \setminus \text{Supp } M_0) \simeq (X_i \setminus \text{Supp } M_i).$$

Since $\text{Supp } M_i = \text{Supp } N_i$ and φ_i is N_i -negative, the image of the p_i -exceptional locus is contained in M_i . Thus we are done if we show that the image of the q_i -exceptional locus is contained in M_{i+1} . Since $X_0 \dashrightarrow X_{i+1}$ does not extract a divisor we have

$$M_{i+1} = (g_i)_* f_i^* M.$$

Yet by the properties above $f_i^* M$ contains the strict transforms of the q_i -exceptional divisors¹⁰. Thus $g_i(\text{Supp } f_i^* M)$ contains the image of the q_i -exceptional locus. By Lemma 6.3 this implies that $\text{Supp}(g_i)_* f_i^* M$ contains the image of the q_i -exceptional locus.

Proof of the claim. For the start of the induction we simply set $V_0 := \Gamma_0$. Indeed since $\text{Supp } M_0 = \text{Supp } N_0$ contains the exceptional locus of the extremal contraction, the divisor $p_0^* M_0$ contains the q_0 -exceptional locus.

For the induction step we make a case distinction: if φ_i is divisorial we simply set $V_i := V_{i-1}$ and note that M_i contains the exceptional divisor since $\text{Supp } M_i = \text{Supp } N_i$. If φ_i is a flip we define V_i as the normalisation of the graph of the bimeromorphic map $V_{i-1} \dashrightarrow \Gamma_i$, and denote by $\alpha_i : V_i \rightarrow V_{i-1}$ and $\beta_i : V_i \rightarrow \Gamma_i$

⁹As usual we define inductively $N_{i+1} := (\varphi_i)_* N_i$. Cp. [Kwc92, 2.26]

¹⁰Note that the q_i -exceptional locus is divisorial since X_{i+1} is $\mathbb{Q}$ -factorial.

the natural maps. We set $f_i := f_{i-1} \circ \alpha_i$ and $g_i := q_i \circ \beta_i$ and summarise the situation in a commutative diagram:

$$\begin{array}{ccccc}
 & & V_i & & \\
 & \swarrow \alpha_i & & \searrow \beta_i & \\
 & V_{i-1} & & \Gamma_i & \\
 f_{i-1} \swarrow & & \searrow g_{i-1} & p_i \swarrow & \searrow q_i \\
 X & & X_i & & X_{i+1}
 \end{array}$$

All we have to show is that the support of f_i^*M contains the g_i -exceptional locus. Note that by our induction hypothesis f_{i-1}^*M contains the g_{i-1} -exceptional locus. Moreover we have

$$M_i = (g_{i-1})_* f_{i-1}^* M$$

and $\text{Supp } M_i = \text{Supp } N_i$. In particular $(g_{i-1})_* f_{i-1}^* M$ contains the image of the p_i -exceptional locus (which is of course equal to the q_i -exceptional locus). Using these two properties the claim follows by elementary set-theoretic computations. $\square$

Proof of Lemma 6.1. Since $\kappa(X) \geq 0$ by [DP03], there exists a $m \in \mathbb{N}$ and an effective Cartier divisor $D \in |mK_X|$. Since $\nu(X) > 0$, the divisor D is not zero. Let

$$\mu : X_0 \rightarrow X$$

be a log-resolution of the divisor D in the following sense: the map μ is an isomorphism on $X \setminus D$, the support of μ^*D has simple normal crossings and X_0 is smooth in a neighbourhood of μ^*D^{11} . Thus if we set $B_0 := (\mu^*D)_{\text{red}}$, the pair (X_0, B_0) is dlt and $X_0 \setminus B_0$ has terminal singularities. Note also that since X has terminal singularities, we have

$$K_{X_0} \sim_{\mathbb{Q}} \mu^* K_X + E$$

with E an effective μ -exceptional $\mathbb{Q}$ -divisor. By construction

$$\text{Supp } E \subset B_0,$$

so the pluricanonical divisor

$$D_0 := \mu^* D + mE \in |mK_{X_0}|$$

satisfies $\text{Supp } D_0 = B_0$. By Remark 6.2 we have

$$\kappa(X) = \kappa(X_0) = \kappa(K_{X_0} + B_0).$$

By Theorem 4.22 there exists a terminating $(K + B)$ -MMP

$$(\varphi_i : (X_i, B_i) \dashrightarrow (X_{i+1}, B_{i+1}))_{i=0, \dots, n}$$

where φ_i is the flip or divisorial contraction of a ray R_i . Then for every $i \in \{1, \dots, n\}$

$$D_{i+1} := \varphi_* D_i \in |mK_{X_{i+1}}|$$

is a pluricanonical divisor such that $\text{Supp } D_{i+1} = B_{i+1}$.

¹¹This last condition can be satisfied since X is a terminal threefold, so it has only finitely many singular points.

For every $0 \leq i \leq n$, if $C \subset X_i$ is a curve with class $[C] \in R_i$, then

$$(K_{X_i} + B_i) \cdot C = \left(\frac{1}{m}D_i + B_i\right) \cdot C < 0,$$

and since $\text{Supp } D_i = B_i$, the curve C is contained in B_i . Thus the exceptional locus of the contraction of the extremal ray R_i is contained in B_i . Now apply Lemma 6.4 to establish an isomorphism

$$\varphi_i|_{X_i \setminus B_i} : (X_i \setminus B_i) \rightarrow (X_{i+1} \setminus B_{i+1}).$$

In conclusion we obtain that $X_{i+1} \setminus B_{i+1}$ has terminal singularities.

The Kodaira dimension of $K + B$ being invariant under a $(K + B)$ -MMP, so Remark 6.2 yields

$$\kappa(X) = \kappa(K_{X_0} + B_0) = \kappa(K_{X_{n+1}} + B_{n+1}).$$

Let now Γ be a desingularisation of the graph of the bimeromorphic map $X_0 \dashrightarrow X_{n+1}$, and denote by $p : \Gamma \rightarrow X_0$ and $q : \Gamma \rightarrow X_{n+1}$ the natural projections. Then the effective divisors

$$(\mu \circ p)^* D \simeq (\mu \circ p)^* mK_X$$

and

$$q^*(D_{n+1} + mB_{n+1}) \simeq mq^*(K_{X_{n+1}} + B_{n+1})$$

are both nef and have the same support. As in [Kwc92, 11.3.3], arguing with a Kähler class, this implies

$$\nu(X) = \nu(K_{X_{n+1}} + B_{n+1}).$$

□

6.5. Lemma. *Let X be a normal compact Kähler space of dimension $n \geq 2$, and let L be a nef $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor. Let $S \subset X$ be an effective $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor such that $L - S$ is nef. Let $T \subset X$ be a prime divisor such that $L|_T \equiv 0$ and $T \not\subset \text{Supp } S$. Then*

$$T \cap \text{Supp } S = \emptyset.$$

Proof. Indeed, if $T \cap \text{Supp } S \neq \emptyset$, then $-S|_T$ is a non-zero antieffective divisor. Thus the restriction $(L - S)|_T \equiv -S|_T$ is not nef, a contradiction. □

6.6. Lemma. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold. Suppose that there exists a $D \in |mK_X|$ with the following properties:*

- *Set $B := \text{Supp } D$. The pair (X, B) is dlt and $X \setminus B$ has terminal singularities.*
- *The divisor $K_X + B$ is nef.*

Let $S \subset B$ be a non-zero effective Weil divisor such that the following holds: for every irreducible component $T \subset B - S$ we have

$$(K_X + B)|_T \equiv 0.$$

Set $X_0 := X$, $D_0 := D$ and $B_0 := B$. Then there exists a finite sequence $K + B - S$ -negative contractions

$$(\varphi_i : (X_i, B_i - S_i) \dashrightarrow (X_{i+1}, B_{i+1} - S_{i+1}))_{i=0, \dots, n}$$

of a ray $R_i \in \overline{NA}(X_i)$ such that the following properties hold:

- a) *The pair $(X_{i+1}, B_{i+1} - S_{i+1})$ is dlt.*
- b) *Set $D_{i+1} := (\varphi_i)_* D_i$. Then the divisor $D_{i+1} \in |mK_{X_{i+1}}|$ satisfies $\text{Supp } D_{i+1} = B_{i+1}$.*

c) We have $(K_{X_i} + B_i) \cdot R_i = 0$ and the divisor $K_{X_{i+1}} + B_{i+1}$ is nef. For every irreducible component $T_{i+1} \subset B_{i+1} - S_{i+1}$ we have

$$(K_{X_{i+1}} + B_{i+1})|_{T_{i+1}} \equiv 0.$$

d) X_{i+1} is $\mathbb{Q}$ -factorial; the pair (X_{i+1}, B_{i+1}) is lc and $X_{i+1} \setminus B_{i+1}$ has terminal singularities.

e) We have $(B_{n+1} - S_{n+1}) \cap S_{n+1} = \emptyset$ and $S_{n+1} \neq \emptyset$.

Remark. Note that we do not claim that $K_{X_{n+1}} + B_{n+1} - S_{n+1}$ is nef, in general this will not be true.

Proof. If $(B - S) \cap S = \emptyset$, there is nothing to prove, so suppose $(B - S) \cap S \neq \emptyset$. Thus $K_X + B - S$ is not nef by Lemma 6.5. Note that $(X_0, B_0 - S_0)$ is dlt.

Step 1. Existence of the MMP. We proceed by induction and assume that X_i, B_i and D_i are already constructed such that properties a), b) and d) hold at level i , moreover $K_{X_i} + B_i$ is nef and $K_{X_i} + B_i|_{T_i} \equiv 0$ for all irreducible components T_i of $B_i - S_i$. If there is no such T_i meeting S_i , then we stop and set $n + 1 = i$. Thus we may assume that $T_i \cap S_i \neq \emptyset$ for some component T_i . We first have to find the extremal ray R_i . Notice that the restriction

$$(K_{X_i} + B_i - S_i)|_{T_i} \equiv -S_i|_{T_i}$$

is not nef, so by Corollary 4.4 there exists a curve $Z_i \subset T_i \subset (B_i - S_i)$ such that

$$(K_{X_i} + B_i - S_i) \cdot Z_i < 0.$$

The inclusion $Z_i \subset (B_i - S_i)$ yields $(K_{X_i} + B_i) \cdot Z_i = 0$ by c). By the cone theorem 1.3 applied to the pair $(X_i, B_i - S_i)$, there exists a decomposition

$$(21) \quad Z_i = \sum \lambda_j C_{i,j} + M_i,$$

where $\lambda_j > 0$, the $C_{i,j}$ are irreducible curves generating a $(K_{X_i} + B_i - S_i)$ -negative extremal in $\overline{\text{NA}}(X_i)$ and $M_i \in \overline{\text{NA}}(X_i)$ such that

$$(K_{X_i} + B_i - S_i) \cdot M_i \geq 0.$$

Let R_i be the extremal ray generated by the curve $C_{i,1}$. Since $K_{X_i} + B_i$ is nef and $(K_{X_i} + B_i) \cdot Z_i = 0$, the decomposition (21) implies that $(K_{X_i} + B_i) \cdot R_i = 0$.

Since $\text{Supp } D_i = B_i$, the $\mathbb{Q}$ -Cartier divisor $K_{X_i} + B_i - S_i$ is $\mathbb{Q}$ -linearly equivalent to an effective divisor with support in B_i . In particular every curve $C \subset X_i$ such that $[C] \in R_i$ is contained in B_i . Since the pair (X_i, B_i) is lc this shows that the support of the extremal ray is contained in an irreducible surface with slc singularities, [Kwc92, 16.9]. Thus the contraction of R_i exists by Theorem 1.4. Moreover, if the ray is small, then the flip exists by Theorem 4.19. We denote by

$$\varphi_i : (X_i, B_i - S_i) \dashrightarrow (X_{i+1}, B_{i+1} - S_{i+1})$$

the flip or divisorial contraction of R_i . As in the proof of Lemma 6.1 we see that the properties a) and b) hold at level $i + 1$.

Since $(K_{X_i} + B_i) \cdot R_i = 0$, Lemma 3.7 proves that $K_{X_{i+1}} + B_{i+1}$ is nef, that the pair (X_{i+1}, B_{i+1}) is lc. and that for every irreducible component $T_{i+1} \subset B_{i+1} - S_{i+1}$, the restriction $(K_{X_{i+1}} + B_{i+1})|_{T_{i+1}} \equiv 0$. Finally, the divisor $K_{X_i} + B_i$ being nef and $\mathbb{Q}$ -linearly equivalent to an effective divisor with support in B_i , Lemma 6.4 implies that $(X_{i+1} \setminus B_{i+1}) \simeq (X_i \setminus B_i)$ has terminal singularities.

Step 2. Termination of the MMP. By Theorem 4.20 we know that after finitely many steps the flipping locus is disjoint from $B_i - S_i$. Yet a $K + B - S$ -negative extremal contraction that is disjoint from $B - S$ is also a K -negative extremal contraction. Thus the sequence terminates by Theorem 4.21. $\square$

6.7. Lemma. *In the situation of Lemma 6.6 suppose additionally that $\nu(K_X + B) = 1$, and let $S \subset B$ be an irreducible component.*

Then there exists a normal $\mathbb{Q}$ -factorial compact Kähler threefold X' that is bimeromorphic to X and has a $D' \in |mK_{X'}|$ with the following properties:

- a) *Set $B' := \text{Supp } D'$. The pair (X', B') is lc and $X' \setminus B'$ has terminal singularities.*
- b) *The divisor $K_{X'} + B'$ is nef with $\nu(K_{X'} + B') = 1$. Moreover, $\kappa(K_X + B) = \kappa(K_{X'} + B')$.*
- c) *There exists an irreducible component $S' \subset B'$ that is a connected component of B' .*
- d) *The pair $(X', B' - S')$ is dlt.*

Proof. We first check that $B - S$ satisfies the positivity condition in Lemma 6.6: indeed if T is any irreducible component of $\text{Supp}(D)$, then $(K_X + B)|_T \equiv 0$, otherwise $\nu(K_X + B) \geq 2$.

Using the notation of Lemma 6.6, set $X' = X_{n+1}$ etc. We know that S_{n+1} is a non-zero irreducible divisor that is disjoint from $B_{n+1} - S_{n+1}$. Thus the property c) holds. Since all the contractions in Lemma 6.6 are $(K + B)$ -trivial, the Kodaira dimension is invariant:

$$\kappa(K_X + B) = \kappa(K_{X_{n+1}} + B_{n+1}).$$

The other properties were already shown in Lemma 6.6. $\square$

6.8. Lemma. *Let X be a normal compact Kähler space with rational singularities of dimension $n \geq 2$. Let D be an effective Cartier divisor. Suppose that D is nef and $\nu(D) \geq 2$. Then the support of D is connected.*

Proof. It is sufficient to prove that $H^1(X, \mathcal{O}_X(-D)) = 0$. Since X has rational singularities we may replace it by a resolution of singularities without changing this cohomology group. Then we have

$$H^1(X, \mathcal{O}_X(-D)) \simeq H^{n-1}(X, K_X \otimes \mathcal{O}_X(D)) = 0$$

by [DP03, Thm.0.1]. $\square$

6.9. Lemma. *In the situation of Lemma 6.6 suppose additionally that $\nu(K_X + B) = 2$.*

Then there exists a normal $\mathbb{Q}$ -factorial compact Kähler threefold X' , bimeromorphic to X such that $(X', 0)$ is klt and a divisor $D' \in |mK_{X'}|$ with the following properties:

- a) *Set $B' := \text{Supp } D'$. The pair (X', B') is lc and $X' \setminus B'$ has terminal singularities.*
- b) *The divisor $K_{X'} + B'$ is nef with $\nu(K_{X'} + B') = 2$. Moreover, $\kappa(K_X + B) = \kappa(K_{X'} + B')$.*
- c) *For every irreducible component $T' \subset B'$ we have $(K_{X'} + B')|_{T'} \not\equiv 0$.*
- d) *$(K_{X'} + B') \cdot K_{X'}^2 \geq 0$.*

Proof. The construction of the Kähler space X' proceeds in two steps.

Part I. Eliminating the $(K + B)$ -trivial components. Let $S \subset B$ be the union of all the irreducible components $T \subset B$ such that $(K_X + B)|_T \neq 0$. Since $\nu(K_X + B) = 2$, the set S is not empty. Thus we can apply Lemma 6.6 to obtain a sequence of $(K + B - S)$ -negative contractions

$$\psi : X \dashrightarrow \hat{X}$$

where $\hat{X}$ is a normal $\mathbb{Q}$ -factorial compact Kähler threefold carrying a divisor $\hat{D} \in |mK_{\hat{X}}|$ subject to the following properties:

- (α) Set $\hat{B} := \text{Supp } \hat{D}$. The pair $(\hat{X}, \hat{B})$ is lc and $\hat{X} \setminus \hat{B}$ has terminal singularities.
- (β) The divisor $K_{\hat{X}} + \hat{B}$ is nef with $\nu(K_{\hat{X}} + \hat{B}) = 2$. Moreover, $\kappa(K_X + B) = \kappa(K_{\hat{X}} + \hat{B})$.
- (γ) Set $\hat{S} := \psi_* S$. Then the pair $(\hat{X}, \hat{B} - \hat{S})$ is dlt.
- (δ) $\text{supp}(\hat{B} - \hat{S}) \cap \hat{S} = \emptyset$ and $\hat{S} \neq 0$.

Notice also that $K_{\hat{X}} + \hat{B}$ is $\mathbb{Q}$ -linearly equivalent to an effective divisor with support in $\hat{B}$. By Lemma 6.8 this implies that $\hat{B} = \hat{S} + (\hat{B} - \hat{S})$ is connected. Thus property (δ) implies that $\hat{B} - \hat{S} = 0$. Hence $(K_{\hat{X}} + \hat{B})|_T$ is non-zero for every irreducible component $T \subset \hat{B}$.

Part II. Eliminating K -negative curves. Setting

$$X_0 := \hat{X}, \quad B_0 := \hat{B}, \quad D_0 := \hat{D},$$

we will next construct a finite sequence of K -negative contractions and flips

$$(\varphi_i : X_i \dashrightarrow X_{i+1})_{i=0, \dots, n}$$

of extremal rays $R_i \in \overline{\text{NA}}(X_i)$ such that the following properties hold:

- (1) X_{i+1} is $\mathbb{Q}$ -factorial and $(X_{i+1}, 0)$ is klt.
- (2) Set $D_{i+1} := (\varphi_i)_* D_i$. Then $D_{i+1} \in |mK_{X_{i+1}}|$ satisfies $\text{Supp } D_{i+1} = B_{i+1}$.
- (3) We have $(K_{X_i} + B_i) \cdot R_i = 0$, the divisor $K_{X_{i+1}} + B_{i+1}$ is nef and $\nu(K_{X_{i+1}} + B_{i+1}) = 2$.
- (4) The pair (X_{i+1}, B_{i+1}) is lc and $X_{i+1} \setminus B_{i+1}$ has terminal singularities.
- (5) For every irreducible component $T \subset B_{i+1}$ we have $(K_{X_{i+1}} + B_{i+1})|_T \neq 0$.
- (6) $(K_{X_{n+1}} + B_{n+1}) \cdot K_{X_{n+1}}^2 \geq 0$.

Note first that $(\hat{X}, 0)$ is klt singularities since $(\hat{X}, \hat{B} - \hat{S}) = (\hat{X}, 0)$ is dlt [KM98, Prop.2.41]. If $(K_{\hat{X}} + \hat{B}) \cdot K_{\hat{X}}^2 \geq 0$, there is nothing to prove, thus we may assume that

$$(K_{\hat{X}} + \hat{B}) \cdot K_{\hat{X}}^2 < 0.$$

We therefore start by showing that $(K_{X_i} + B_i) \cdot K_{X_i}^2 < 0$ implies the existence of a K_{X_i} -negative contraction that is $(K_{X_i} + B_i)$ -trivial. Let $B_i = \sum B_{i,l}$ be the decomposition of B_i in its irreducible components. Since the pair (X_i, B_i) is lc, the surfaces $B_{i,l}$ have slc singularities. Moreover, by adjunction [Kwc92, Prop.16.9], there exists a boundary divisor $\Delta_{i,l}$ such that

$$K_{B_{i,l}} + \Delta_{i,l} \sim_{\mathbb{Q}} (K_{X_i} + B_i)|_{B_{i,l}},$$

and the pair $(B_{i,l}, \Delta_{i,l})$ is slc. By Proposition 5.3 this implies that

$$(K_{X_i} + B_i)|_{B_{i,l}} \sim_{\mathbb{Q}} Z_{i,l}$$

with $Z_{i,l}$ an effective 1-cycle.

Suppose that $K_{X_i}|_{B_{i,l}} \cdot Z_{i,l} \geq 0$ for all l . Since

$$K_{X_i} \sim_{\mathbb{Q}} \sum a_{i,l} B_{i,l}$$

with $a_{i,l} > 0$ we conclude

$$(K_{X_i} + B_i) \cdot K_{X_i}^2 = \sum a_{i,l} (K_{X_i} + B_i) \cdot K_{X_i} \cdot B_{i,l} = \sum a_{i,l} K_{X_i}|_{B_{i,l}} \cdot Z_{i,l} \geq 0,$$

a contradiction. Thus we can suppose (up to renumbering) that

$$K_{X_i}|_{B_{i,1}} \cdot Z_{i,1} < 0.$$

By the cone theorem 1.3 applied to the pair $(X_i, 0)$, there exists a decomposition

$$(22) \quad Z_{i,1} = \sum \lambda_j C_{i,j} + M_i,$$

where $\lambda_j > 0$, the $C_{i,j}$ are irreducible curves generating a K_{X_i} -negative extremal in $\overline{\text{NA}}(X_i)$ and where $M_i \in \overline{\text{NA}}(X_i)$ is a class satisfying $K_{X_i} \cdot M_i \geq 0$.

Notice now the following: since $K_{X_i} + B_i$ is $\mathbb{Q}$ -linearly equivalent to an effective divisor with support B_i and $(K_{X_i} + B_i)^3 = 0$ we have

$$(K_{X_i} + B_i)^2 \cdot B_{i,l} = 0$$

for all l . In particular $(K_{X_i} + B_i) \cdot Z_{i,1} = 0$. Since $K_{X_i} + B_i$ is nef we deduce from (22) that

$$(23) \quad (K_{X_i} + B_i) \cdot C_{i,1} = 0.$$

Let now R_i be the extremal ray generated by $C_{i,1}$. As in the proof of Lemma 6.6, the locus of R_i is contained in a surface with slc singularities (a component of B_i). Thus the contraction of R_i exists by Theorem 1.4 and, if the extremal ray is small, the flip exists by Theorem 4.19. We denote by

$$\varphi_i : (X_i, 0) \dashrightarrow (X_{i+1}, 0)$$

the flip or divisorial contraction of R_i . As in the proof of Lemma 6.1 we see that the properties (1) and (2) hold at level $i + 1$.

By the induction hypothesis, the divisor $K_{X_i} + B_i$ is nef of numerical dimension two and by (23) numerically trivial on the extremal ray R_i . Using Lemma 3.7 this implies that $K_{X_{i+1}} + B_{i+1}$ is nef of numerical dimension two, moreover the pair (X_{i+1}, B_{i+1}) is lc. Finally Lemma 6.4 implies that $(X_{i+1} \setminus B_{i+1}) \simeq (X_i \setminus B_i)$ has terminal singularities.

Part III In total we have constructed a sequence of K -negative contractions satisfying the properties (1)-(5). By Theorem 4.21 any such sequence terminates after n steps. By construction this yields

$$(K_{X_{n+1}} + B_{n+1}) \cdot K_{X_{n+1}}^2 \geq 0.$$

Since all the contractions are $K + B$ -trivial, the Kodaira dimension is invariant, i.e. we have $\kappa(K_{\hat{X}} + \hat{B}) = \kappa(K_{X_{n+1}} + B_{n+1})$. $\square$

7. POSITIVITY OF COTANGENT SHEAVES

In this section we briefly review stability and Chern classes on singular complex spaces, then we prove the crucial Chern class inequality Theorem 7.11. For simplicity of notations we restrict ourselves to the case of threefolds with isolated singularities - which is all we need later - but all the statements can easily be adapted for spaces of arbitrary dimension that are smooth in codimension two.

7.1. Definition. *Let X be a normal compact Kähler threefold with isolated singularities. Let $\mathcal{F}_1$ and $\mathcal{F}_2$ be coherent sheaves on X , and let $\pi : \hat{X} \rightarrow X$ be a log-resolution.*

Then the Chern classes $c_i(\pi^\mathcal{F}_1)$ and $c_i(\pi^*\mathcal{F}_2)$ are well-defined elements of $H^{2i}(\hat{X}, \mathbb{Z})$ [TT86], see also [Gri10]. Thus for every $\alpha \in N^1(X)$ the intersection numbers*

$$\pi^*\alpha \cdot c_1(\pi^*\mathcal{F}_1) \cdot c_1(\pi^*\mathcal{F}_2) \quad \text{and} \quad \pi^*\alpha \cdot c_2(\pi^*\mathcal{F}_1)$$

are well-defined, so by the duality $N_1(X) = N^1(X)^$ we define*

$$c_1(\mathcal{F}_1) \cdot c_1(\mathcal{F}_2) \in N_1(X) : \alpha \mapsto \pi^*\alpha \cdot c_1(\pi^*\mathcal{F}_1) \cdot c_1(\pi^*\mathcal{F}_2)$$

and

$$c_2(\mathcal{F}_1) \in N_1(X) : \alpha \mapsto \pi^*\alpha \cdot c_2(\pi^*\mathcal{F}_1).$$

7.2. Lemma. *In the situation of Definition 7.1, the classes $c_1(\mathcal{F}_1) \cdot c_1(\mathcal{F}_2)$ and $c_2(\mathcal{F}_1)$ do not depend on the choice of the desingularisation. Moreover if*

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{Q} \rightarrow 0$$

is an exact sequence of coherent sheaves, then we have the usual formula

$$c_2(\mathcal{G}) = c_2(\mathcal{F}) + c_2(\mathcal{Q}) + c_1(\mathcal{F}) \cdot c_1(\mathcal{Q}).$$

Proof. In order to see for instance that the definition $c_2(\mathcal{F})$ does not depend on the resolution it suffices to consider the case where $\pi_1 : \hat{X}_1 \rightarrow X$ and $\pi_2 : \hat{X}_2 \rightarrow X$ are two log-resolutions with a factorisation $q : X_2 \rightarrow X_1$. Then we want to show that

$$(q^*\pi_1^*\alpha) \cdot c_2(q^*\pi_1^*\mathcal{F}) = (\pi_1^*\alpha) \cdot c_2(\pi_1^*\mathcal{F}).$$

This follows from the general fact, that, given a holomorphic map $f : X \rightarrow Y$ between compact complex manifolds and $\mathcal{G}$ a coherent sheaf on Y , then $c_j(f^*(\mathcal{G})) = f^*(c_j(\mathcal{G}))$, cp. e.g. [Gri10].

Next consider a log-resolution $\pi : \hat{X} \rightarrow X$ with exceptional locus $D = \sum D_k$ and $\mathcal{F}$ a coherent sheaf with support on D . Let $i^k : D_k \rightarrow X$ be the inclusion. Then by the Grothendieck-Riemann-Roch formula, see e.g., [Gri10, Thm1.1] (and [Ful98, Thm.15.2] in the algebraic case),

$$\text{ch}(i_*^k(\mathcal{F}|_{D_k})) = i_*^k(\text{td}(N_{D_k/X})^{-1} \cdot \text{ch}(\mathcal{F}|_{D_k})).$$

Thus if $\alpha \in N^1(X)$, then the projection formula gives

$$\pi^*\alpha \cdot c_2(i_*^k(\mathcal{F}|_{D_k})) = 0.$$

Now an easy induction on the number of components of D shows that

$$(24) \quad \pi^*(\alpha) \cdot c_2(\mathcal{F}) = 0$$

For the second statement we consider the pull-back

$$\pi^*\mathcal{F} \xrightarrow{\alpha} \pi^*\mathcal{G} \rightarrow \pi^*\mathcal{Q} \rightarrow 0$$

by a log-resolution $\pi : \hat{X} \rightarrow X$. In general the morphism α is not injective, but its kernel has support in the π -exceptional locus. Using the usual rules for computing Chern classes on compact manifolds and invoking (24), we have

$$\pi^* \alpha \cdot c_2(\pi^* \mathcal{F}) = \pi^* \alpha \cdot c_2(\pi^* \mathcal{F} / \ker \alpha)$$

and

$$\pi^* \alpha \cdot c_1(\pi^* \mathcal{F}) \cdot c_1(\pi^* \mathcal{Q}) = \pi^* \alpha \cdot c_1(\pi^* \mathcal{F} / \ker \alpha) \cdot c_1(\pi^* \mathcal{Q}).$$

Thus the statement follows from the standard formula in the smooth case. $\square$

7.3. Remark. In the proof of Lemma 7.2 we have shown the following. Let $\pi : \hat{X} \rightarrow X$ be a log resolution of the Kähler threefold X with only isolated singularities. Let $\mathcal{F}$ be a coherent sheaf on $\hat{X}$, supported on the exceptional divisor D of π and let $\alpha \in N^1(X)$. Then we have

$$\pi^*(\alpha) \cdot c_2(\mathcal{F}) = \pi^*(\alpha) \cdot c_1(\mathcal{F})^2 = 0.$$

Given a normal compact Kähler threefold X with isolated singularities and a torsion-free sheaf $\mathcal{F}$ there is no obvious candidate for the first Chern class $c_1(\mathcal{F}) \in H^2(X, \mathbb{R})$. However we can define, as in the situation of Definition 7.1, for every $\alpha \in N^1(X)$ the intersection number $\alpha^2 \cdot c_1(\mathcal{F})$ by pulling-back to some log-resolution $\pi : \hat{X} \rightarrow X$:

$$\alpha^2 \cdot c_1(\mathcal{F}) := (\pi^*(\alpha))^2 \cdot c_1(\pi^*(\mathcal{F})).$$

7.4. Definition. Let X be a normal compact Kähler threefold with isolated singularities and let α be a nef class on X . We say that a non-zero torsion-free sheaf $\mathcal{F}$ is α -semistable (resp. α -stable) if for every non-zero saturated subsheaf $\mathcal{E} \subset \mathcal{F}$ we have

$$\mu_\alpha(\mathcal{E}) := \frac{\alpha^2 \cdot c_1(\mathcal{E})}{\text{rk} \mathcal{E}} \leq \frac{\alpha^2 \cdot c_1(\mathcal{F})}{\text{rk} \mathcal{F}} =: \mu_\alpha(\mathcal{F}) \quad (\text{resp. } <).$$

7.5. Proposition. (Harder-Narasimhan filtration) Let X be a normal compact Kähler threefold with isolated singularities, and let α be a nef class on X . Let $\mathcal{F}$ be a non-zero torsion-free coherent sheaf on X . Then there exists a filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_k = \mathcal{F}$$

such that for every $i \in \{1, \dots, k\}$ the quotient $\mathcal{F}_i / \mathcal{F}_{i-1}$ is α -semistable and we have a strictly decreasing sequence of slopes

$$\mu_\alpha(\mathcal{F}_i / \mathcal{F}_{i-1}) > \mu_\alpha(\mathcal{F}_{i+1} / \mathcal{F}_i) \quad \forall i \in 1, \dots, k-1.$$

Proof. We proceed as in [Kob87]; the main point is to show that there is a constant C such that

$$\mu_\alpha(\mathcal{F}) \leq C$$

for all $\mathcal{F} \subset \mathcal{E}$. First we reduce to the smooth case: take a log resolution $\pi : \hat{X} \rightarrow X$ such that $\pi^*(\mathcal{E})^{**}$ is locally free and observe that $\mu_\alpha(\mathcal{F}) = \mu_{\pi^*(\alpha)}(\pi^*(\mathcal{F}) / \text{tor})$.

So we may assume X smooth and $\mathcal{E}$ locally free. Then the proof of [Kob87, Lemma 7.16] works. Once the boundedness is settled, the existence of the filtration is shown as in the classical case. $\square$

We shall also use the following elementary result, cp. [Kob87, Thm.7.18].

7.6. Proposition. (*Jordan-Hölder filtration*) Let X be a normal compact Kähler threefold with isolated singularities, and let α be a nef class on X . Let $\mathcal{F}$ be a non-zero α -semi-stable torsion-free coherent sheaf on X . Then there exists a filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_k = \mathcal{F}$$

such that for every $i \in \{1, \dots, k\}$ the quotient $\mathcal{F}_i/\mathcal{F}_{i-1}$ is α -stable.

7.7. Definition. Let X be a normal compact Kähler threefold, and let α be a nef class on X . A non-zero torsion-free coherent sheaf $\mathcal{F}$ on X is α -generically nef if for every torsion-free quotient sheaf $\mathcal{F} \rightarrow Q \rightarrow 0$ we have

$$\alpha^2 \cdot c_1(Q) \geq 0.$$

7.8. Remark. In the situation of Definition 7.7, let

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_k = \mathcal{F}$$

be the Harder-Narasimhan filtration with respect to α . If $\mathcal{F}$ is α -generically nef, then by definition $\alpha^2 \cdot c_1(\mathcal{F}/\mathcal{F}_{k-1}) \geq 0$. Thus

$$\alpha^2 \cdot c_1(\mathcal{F}_i/\mathcal{F}_{i-1}) \geq 0$$

for every $i \in \{1, \dots, k\}$.

We can now prove the Bogomolov inequality for *stable* sheaves on a singular space.

7.9. Theorem. Let X be a normal compact Kähler threefold with isolated singularities, and let α be a Kähler class on X . Let $\mathcal{F}$ be an α -stable non-zero torsion-free coherent sheaf on X . Then we have

$$\alpha \cdot c_2(\mathcal{F}) \geq \left(\frac{r-1}{2r}\right) \alpha \cdot c_1^2(\mathcal{F}).$$

Proof. We fix a log-resolution $\pi : \hat{X} \rightarrow X$.

Step 1. Suppose that $\mathcal{F}$ is reflexive. Set $\hat{\mathcal{F}} := (\pi^*\mathcal{F})^{**}$. Since $\mathcal{F}$ is reflexive, the sheaves $\hat{\mathcal{F}}$ and $\pi^*\mathcal{F}$ coincide in the complement of the π -exceptional locus. In particular by Remark 7.3 one has

$$\alpha \cdot c_2(\mathcal{F}) = \pi^*\alpha \cdot c_2(\pi^*\mathcal{F}) = \pi^*\alpha \cdot c_2(\hat{\mathcal{F}})$$

and

$$\alpha \cdot c_1^2(\mathcal{F}) = \pi^*\alpha \cdot c_1^2(\pi^*\mathcal{F}) = \pi^*\alpha \cdot c_1^2(\hat{\mathcal{F}}).$$

Thus it is sufficient to prove the inequality for $\hat{\mathcal{F}}$. Arguing exactly as in the proof of [DP03, Prop.6.9] we see that $\hat{\mathcal{F}}$ is $\pi^*\alpha$ -stable. Since stability is an open property [Cao13, Prop.2.1] we obtain that $\hat{\mathcal{F}}$ is $(\pi^*\alpha + \varepsilon\omega)$ -stable where ω is a Kähler form on $\hat{X}$ and $0 < \varepsilon \ll 1$. Thus [BS94] yields

$$(\pi^*\alpha + \varepsilon\omega) \cdot c_2(\hat{\mathcal{F}}) \geq \frac{r-1}{2r} (\pi^*\alpha + \varepsilon\omega) \cdot c_1^2(\hat{\mathcal{F}})$$

for every $0 < \varepsilon \ll 1$. The claim follows by passing to the limit $\varepsilon \rightarrow 0$.

Step 2. Reduction to the case where $\mathcal{F}$ is reflexive. Since $\mathcal{F}$ is torsion-free, the injection $i : \mathcal{F} \hookrightarrow \mathcal{F}^{**}$ is an isomorphism in codimension one. Thus the kernel and cokernel of

$$\pi^*(\mathcal{F}) \rightarrow \pi^*(\mathcal{F}^{**})$$

have support in the union of the π -exceptional locus and a set of dimension at most one. In particular $c_1(\pi^*\mathcal{F}) = c_1(\pi^*\mathcal{F}^{**}) + D$ with D a π -exceptional divisor, so the projection formula yields

$$\alpha \cdot c_1^2(\mathcal{F}) = \alpha \cdot c_1^2(\mathcal{F}^{**}).$$

Thus we are done if we prove that $\alpha \cdot c_2(\mathcal{F}^{**}) \geq \alpha \cdot c_2(\mathcal{F})$. By the second part of Lemma 7.2 it is sufficient to prove that

$$\alpha \cdot c_2(\mathcal{F}^{**}/\mathcal{F}) = \pi^*(\alpha) \cdot c_2(\pi^*(\mathcal{F}^{**}/\mathcal{F})) \geq 0.$$

Let S be the union of the 1-dimensional irreducible components of the support of $\mathcal{F}^{**}/\mathcal{F}$ and $\hat{S}$ the strict transform in $\hat{X}$; we may assume that the irreducible components $\hat{S}_i$ of $\hat{S}$ are smooth. Set

$$Q := (i_{\hat{S}})_*(\pi^*(\mathcal{F}^{**}/\mathcal{F})|_{\hat{S}}).$$

Then by Remark 7.3 and Lemma 7.2, it suffices to show

$$\pi^*(\alpha) \cdot c_2(Q) \geq 0.$$

Yet Q has support on a set of dimension one, so the Grothendieck Riemann-Roch formula yields

$$c_2(Q) = \sum a_i \hat{S}_i,$$

where $a_i \in \mathbb{N}_0$. Since α is nef, the statement follows. $\square$

7.10. Lemma. *Let X be a compact Kähler threefold with isolated singularities. Let α be a nef class on X , and let $\mathcal{F}$ be a coherent sheaf. Then we have*

$$(\alpha^2 \cdot c_1(\mathcal{F}))^2 \geq (\alpha \cdot c_1^2(\mathcal{F})) \cdot \alpha^3.$$

Proof. By our definition of the intersection numbers 7.1 we may suppose X smooth. Since α is a limit of Kähler classes, it is sufficient to prove the statement under the stronger hypothesis that α is a Kähler class. Now the inequality follows from the usual Hodge index theorem [Voi02, Thm.6.33] by an elementary computation. $\square$

7.11. Theorem. *Let (X, ω) be a compact Kähler threefold with isolated singularities, and let $\mathcal{F}$ be a non-zero reflexive coherent sheaf on X such that $\det \mathcal{F}$ is $\mathbb{Q}$ -Cartier. Suppose that there exists a pseudoeffective class $P \in N^1(X)$ such that*

$$L := c_1(\mathcal{F}) + P$$

is a nef class. Suppose furthermore that for all $0 < \varepsilon \ll 1$ the sheaf $\mathcal{F}$ is $(L + \varepsilon\omega)$ -generically nef. Then we have

$$L \cdot c_2(\mathcal{F}) \geq \frac{1}{2}(L \cdot c_1^2(\mathcal{F}) - L^3).$$

In particular, if $L \cdot c_1^2(\mathcal{F}) \geq 0$ and $L^3 = 0$, then

$$(25) \quad L \cdot c_2(\mathcal{F}) \geq 0.$$

Remark. This statement (and its proof) is a variation of [Miy87, Thm.6.1], [Kwc92, Prop.10.12]. However the weaker assumptions will be crucial for the application in the proof of Theorem 8.2.

Proof. Fix $0 < \varepsilon \ll 1$, and consider the Harder-Narasimhan filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_k = \mathcal{F}$$

for $\mathcal{F}$ with respect to $L_\varepsilon := L + \varepsilon\omega$. Then by Lemma 7.2

$$(26) \quad L_\varepsilon \cdot c_2(\mathcal{F}) = L_\varepsilon \cdot \left(\sum_{1 \leq i < j \leq k} c_1(\mathcal{F}_i/\mathcal{F}_{i-1})c_1(\mathcal{F}_j/\mathcal{F}_{j-1}) + \sum_{i=1}^k c_2(\mathcal{F}_i/\mathcal{F}_{i-1}) \right).$$

Since $\mathcal{F}$ is L_ε -generically nef, by Remark 7.8

$$(27) \quad L_\varepsilon^2 \cdot c_1(\mathcal{F}_i/\mathcal{F}_{i-1}) \geq 0 \quad \forall i \in \{1, \dots, k\}.$$

For every $i \in \{1, \dots, k\}$, let

$$0 = \mathcal{F}_{i,0} \subset \mathcal{F}_{i,1} \subset \dots \subset \mathcal{F}_{i,k_i} = \mathcal{F}_i/\mathcal{F}_{i-1}$$

be the Jordan-Hölder filtration of $\mathcal{F}_i/\mathcal{F}_{i-1}$ with respect to L_ε . Then we have

$$(28) \quad \begin{aligned} L_\varepsilon \cdot c_2(\mathcal{F}_i/\mathcal{F}_{i-1}) &= \\ &= L_\varepsilon \cdot \left(\sum_{1 \leq p < q \leq k_i} c_1(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1})c_1(\mathcal{F}_{i,q}/\mathcal{F}_{i,q-1}) + \sum_{p=1}^{k_i} c_2(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}) \right). \end{aligned}$$

Since $\mathcal{F}_i/\mathcal{F}_{i-1}$ is L_ε -semistable with non-negative slope by (27), we obtain

$$(29) \quad L_\varepsilon^2 \cdot c_1(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}) \geq 0 \quad \forall i \in \{1, \dots, k\}, p \in \{1, \dots, k_i\}.$$

Plugging the equations (28) into the equation (26) and following the lexicographic order, we rename the graded pieces $\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}$ into $\mathcal{G}_l$, where $l \in \{1, \dots, n\}$. Thus

$$(30) \quad L_\varepsilon \cdot c_2(\mathcal{F}) = L_\varepsilon \cdot \left(\sum_{1 \leq l < m \leq n} c_1(\mathcal{G}_l)c_1(\mathcal{G}_m) + \sum_{l=1}^n c_2(\mathcal{G}_l) \right),$$

where the $\mathcal{G}_l$ are non-zero torsion-free L_ε -stable sheaves. Moreover, by (29) we have

$$(31) \quad L_\varepsilon^2 \cdot c_1(\mathcal{G}_l) \geq 0 \quad \forall l \in \{1, \dots, n\}.$$

Since we renamed according to the lexicographic order, the sequence of slopes with respect to L_ε is (not necessarily strictly) decreasing. Thus, setting $r_l := \text{rk} \mathcal{G}_l$ and

$$\alpha_l := \frac{L_\varepsilon^2 \cdot c_1(\mathcal{G}_l)}{r_l L_\varepsilon^3},$$

we obtain

$$(32) \quad \alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n \geq 0.$$

By Theorem 7.9 we have $L_\varepsilon \cdot c_2(\mathcal{G}_l) \geq \left(\frac{r_l-1}{2r_l}\right) L_\varepsilon \cdot c_1^2(\mathcal{G}_l)$ for all $l \in \{1, \dots, n\}$. Therefore using (30) we obtain

$$\begin{aligned} L_\varepsilon \cdot c_2(\mathcal{F}) &= L_\varepsilon \cdot \left(\frac{1}{2} c_1(\mathcal{F})^2 + \sum_{l=1}^n c_2(\mathcal{G}_l) - \frac{1}{2} \sum_{l=1}^n c_1(\mathcal{G}_l)^2 \right) \\ &\geq L_\varepsilon \cdot \left(\frac{1}{2} c_1(\mathcal{F})^2 - \frac{1}{2r_l} \sum_{l=1}^n c_1(\mathcal{G}_l)^2 \right). \end{aligned}$$

By Lemma 7.10 we have

$$(L_\varepsilon^2 \cdot c_1(\mathcal{G}_l))^2 \geq (L_\varepsilon \cdot c_1^2(\mathcal{G}_l)) \cdot L_\varepsilon^3 \quad \forall l \in \{1, \dots, n\},$$

so, using the coefficient α_l defined above, we get $L_\varepsilon \cdot c_1^2(\mathcal{G}_i) \leq \alpha_i^2 r_i^2 L_\varepsilon^3$. Putting this into the last inequality yields

$$(33) \quad L_\varepsilon \cdot c_2(\mathcal{F}) \geq L_\varepsilon \cdot \left(\frac{1}{2} c_1(\mathcal{F})^2 - \frac{1}{2} \sum_{l=1}^n \alpha_l^2 r_l L_\varepsilon^2 \right)$$

$$(34) \quad = \frac{1}{2} L_\varepsilon \cdot \left((c_1(\mathcal{F})^2 - L_\varepsilon^2) + (1 - \sum_{l=1}^n \alpha_l^2 r_l) L_\varepsilon^2 \right)$$

We claim that

$$(1 - \sum_{l=1}^n \alpha_l^2 r_l) \geq 0.$$

Assuming this for the time being, let us see how to conclude : since $L_\varepsilon^3 > 0$, the claim together with (34) yields

$$L_\varepsilon \cdot c_2(\mathcal{F}) \geq \frac{1}{2} L_\varepsilon \cdot (c_1(\mathcal{F})^2 - L_\varepsilon^2).$$

Now we take the limit $\varepsilon \rightarrow 0$.

Proof of the claim. First of all, (32) implies that

$$1 - \sum_{l=1}^n \alpha_l^2 r_l \geq 1 - \alpha_1 \sum_{l=1}^n r_l \alpha_l.$$

Now by definition of the numbers α_l we have

$$\sum_{l=1}^n r_l \alpha_l = \sum_{l=1}^n \frac{L_\varepsilon^2 \cdot c_1(\mathcal{G}_l)}{L_\varepsilon^3} = \frac{L_\varepsilon^2 \cdot c_1(\mathcal{F})}{L_\varepsilon^3}.$$

Yet by definition of L and L_ε ,

$$c_1(\mathcal{F}) = L - P = L_\varepsilon - P - \varepsilon \omega.$$

Since P is pseudoeffective and since ω is Kähler it follows that $L_\varepsilon^2 \cdot c_1(\mathcal{F}) \leq L_\varepsilon^3$, hence $\sum_{l=1}^n r_l \alpha_l \leq 1$. Since $r_l \geq 1$ and $\alpha_l \geq 0$ for all $l \in \{1, \dots, n\}$ this implies $\alpha_1 \leq 1$, proving the claim. $\square$

8. ABUNDANCE

In this section, we establish the abundance theorem for non-algebraic Kähler threefolds. The main difficulty is to show that if the numerical Kodaira dimension $\nu(X) = 1$ or $\nu(X) = 2$, then $\kappa(X) \geq 1$. This will be done in Theorems 8.1 and 8.2.

8.A. The case $\nu = 1$.

8.1. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. If $\nu(X) = 1$, then $\kappa(X) \geq 1$.*

Proof. By Lemma 6.1 and Lemma 6.7 we are reduced to proving the following statement:

Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold such that $(X, 0)$ is klt, carrying a divisor $D \in |mK_X|$ with the following properties:

- a) Set $B := \text{Supp } D$. The pair (X, B) is lc and $X \setminus B$ has terminal singularities.*
- b) The divisor $K_X + B$ is nef and we have $\nu(K_X + B) = 1$.*
- c) There exists an irreducible component $S \subset B$ that is a connected component.*

Then $\kappa(X) \geq 1$.

We follow the argument in [Kwc92, Ch.13]: by adjunction [Kwc92, 16.9.1] there exists a boundary divisor Δ on S such that (S, Δ) is slc and $K_S + \Delta$ is numerically trivial. By Proposition 5.1 the divisor $K_S + \Delta$ is torsion. Due to a covering trick of Miyaoka [Kwc92, 11.3.6] we may suppose after a finite cover, étale in codimension one, that

$$\omega_S = \mathcal{O}_S(K_S) \simeq \mathcal{O}_S \simeq \mathcal{O}_S(S).$$

Note that by [KM98, Prop.5.20(4)] the klt property is preserved under a finite morphism which is étale in codimension one. In particular, X is Cohen-Macaulay. By [Kwc92, 11.3.7] we are finished if we prove that for every infinitesimal neighbourhood S_n , the restriction

$$H^p(S_n, \mathcal{O}_{S_n}) \rightarrow H^p(S, \mathcal{O}_S)$$

is surjective for every $p \in \mathbb{N}$. Observe here that condition (3) in [Kwc92, 11.3.7] is satisfied by [Kwc92, 12.1.2], as explained in [Kwc92, p.158]. In fact, using the commutative diagram

$$\begin{array}{ccc} H^p(S_n, \mathbb{C}) & \longrightarrow & H^p(S_n, \mathcal{O}_{S_n}) \\ \downarrow & & \downarrow \\ H^p(S, \mathbb{C}) & \longrightarrow & H^p(S, \mathcal{O}_S) \end{array}$$

and the isomorphism $H^p(S_n, \mathbb{C}) \simeq H^p(S, \mathbb{C})$ we see that it is sufficient to prove that

$$H^p(S, \mathbb{C}) \rightarrow H^p(S, \mathcal{O}_S)$$

is surjective. This is done in Theorem 5.4. $\square$

8.B. The case $\nu = 2$.

8.2. Theorem. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. If $\nu(X) = 2$, then $\kappa(X) \geq 1$.*

The basic idea of the proof is the same as in the projective case [Kwc92, Sect.14], however the computations get considerably simplified by our generalisation of Miyaoka's Chern class inequality [Miy87, Thm.6.1]. Let us start by recalling the Riemann-Roch formula for terminal threefolds:

8.3. Proposition. *Let X be a normal compact Kähler threefold with at most terminal singularities, and let L be a line bundle on X . Then we have*

$$\chi(X, L) = \frac{L^3}{6} - \frac{1}{4}K_X \cdot L^2 + L \cdot \frac{K_X^2 + c_2(X)}{12} + \chi(X, \mathcal{O}_X),$$

where $L \cdot c_2(X) := \pi^*L \cdot c_2(\hat{X})$ with $\pi : \hat{X} \rightarrow X$ any resolution of singularities (cf. Definition 7.1).

Proof. Let $\pi : \hat{X} \rightarrow X$ be a resolution of singularities, which is an isomorphism over the smooth locus of X . Since X has rational singularities, we have $\chi(X, L) = \chi(\hat{X}, \pi^*L)$. Riemann-Roch on the smooth Kähler threefold $\hat{X}$ yields

$$(35) \quad \chi(\hat{X}, \pi^*L) = \frac{\pi^*L^3}{6} - \frac{1}{4}K_{\hat{X}} \cdot (\pi^*L)^2 + \pi^*L \cdot \frac{K_{\hat{X}}^2 + c_2(\hat{X})}{12} + \chi(\hat{X}, \mathcal{O}_{\hat{X}}).$$

Now $\pi^*L^3 = L^3$, and, using again the rationality of the singularities of X , $\chi(\hat{X}, \mathcal{O}_{\hat{X}}) = \chi(X, \mathcal{O}_X)$. Since X is smooth in codimension two we may write

$$K_{\hat{X}} \sim_{\mathbb{Q}} \pi^*K_X + E$$

with E a divisor such that $\pi(E)$ is finite. In particular the projection formula gives $K_{\hat{X}} \cdot (\pi^*L)^2 = K_X \cdot L^2$ and $\pi^*L \cdot K_{\hat{X}}^2 = L \cdot K_X^2$. Thus (35) gives our claim. $\square$

We will also need the following Kähler version of Miyaoka's generic nefness theorem, due to Enoki in the smooth case:

8.4. Proposition. *Let X be a normal compact Kähler space of dimension n with canonical singularities. Suppose that K_X is nef or $\kappa(X) \geq 0$. Then Ω_X is generically nef with respect to any nef class α , i.e. for every torsion-free quotient sheaf*

$$\Omega_X \rightarrow \mathcal{Q} \rightarrow 0,$$

we have $\alpha^{n-1} \cdot c_1(\mathcal{Q}) \geq 0$.

Proof. Fix a nef class $\alpha \in N^1(X)$, and let $\Omega_X \rightarrow \mathcal{Q} \rightarrow 0$ be a torsion-free quotient sheaf. Let $\pi : \hat{X} \rightarrow X$ be a desingularisation by a compact Kähler manifold, and let $\mathcal{K}$ be the kernel of the induced epimorphism

$$(\pi^*\Omega_X)/\text{torsion} \rightarrow (\pi^*\mathcal{Q})/\text{torsion} \rightarrow 0.$$

Using the injective map

$$(\pi^*\Omega_X)/\text{torsion} \hookrightarrow \Omega_{\hat{X}}$$

we may view $\mathcal{K}$ as a subsheaf of $\Omega_{\hat{X}}$, and we denote by $\hat{\mathcal{K}}$ its saturation in $\Omega_{\hat{X}}$. Set

$$\hat{\mathcal{Q}} := \Omega_{\hat{X}}/\hat{\mathcal{K}},$$

then $\hat{\mathcal{Q}}$ is a torsion-free quotient of $\Omega_{\hat{X}}$ coinciding with $(\pi^*\mathcal{Q})/\text{torsion}$ in the complement of the exceptional locus. In particular we have

$$\alpha^{n-1} \cdot c_1(\mathcal{Q}) = (\pi^*\alpha)^{n-1} \cdot c_1((\pi^*\mathcal{Q})/\text{torsion}) = (\pi^*\alpha)^{n-1} \cdot c_1(\hat{\mathcal{Q}}).$$

We will now verify the conditions of [Eno88, Thm.1.4] in order to conclude that $(\pi^*\alpha)^{n-1} \cdot c_1(\hat{\mathcal{Q}}) \geq 0$: since X has canonical singularities we have

$$K_{\hat{X}} = \pi^*K_X + E,$$

with E an effective, π -exceptional $\mathbb{Q}$ -divisor.

Thus if K_X is nef the conditions of [Eno88, Thm.1.4] hold by setting $L = \pi^*K_X$ and $D = E$. If $\kappa(X) \geq 0$ we have $\pi^*K_X = F$ with F an effective $\mathbb{Q}$ -divisor, so the conditions are satisfied by setting $L = 0$ and $D = F + E$. In both cases Enoki's theorem tells us that

$$\omega^{n-1} \cdot c_1(\hat{\mathcal{Q}}) \geq 0$$

for every Kähler form ω on $\hat{X}$. Since $\pi^*\alpha$ is nef, the statement follows by passing to the limit. $\square$

We will use the following Serre vanishing property:

8.5. Lemma. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold, and let $B_1, \dots, B_k$ be prime Weil divisor on X such that B_i is Cohen-Macaulay for every $i \in \{1, \dots, k\}$. Let L be a nef Cartier divisor on X such that $L|_{B_i} \not\equiv 0$ for every $i \in \{1, \dots, k\}$. Let $Y \subset X$ be a subscheme such that $Y_{\text{red}} \subset \sum_{i=1}^k B_i$ and let $\mathcal{F}$ be a coherent sheaf on Y . Then there exists a number $n_0 \in \mathbb{N}$ such that*

$$H^2(Y, \mathcal{F} \otimes \mathcal{O}_Y(L^{\otimes n})) = 0$$

for every $n \geq n_0$.

Proof. Note first that we may suppose that Y is defined by an ideal sheaf $\mathcal{O}_X(-\sum_{i=1}^k a_i B_i)$ with $a_i \in \mathbb{N}$. Indeed, at the general point of every surface B_i , the scheme Y is isomorphic to a scheme $\sum_{i=1}^k a_i B_i$ defined by the ideal sheaf $\mathcal{O}_X(-\sum_{i=1}^k a_i B_i)$. Thus if we consider the restriction map

$$\mathcal{F} \rightarrow \mathcal{F} \otimes \mathcal{O}_{\sum_{i=1}^k a_i B_i},$$

its kernel has support on a scheme of dimension at most one. Hence

$$H^2(Y, \mathcal{F} \otimes \mathcal{O}_Y(L^{\otimes n})) \simeq H^2\left(\sum_{i=1}^k a_i B_i, \mathcal{F} \otimes \mathcal{O}_{\sum_{i=1}^k a_i B_i}(L^{\otimes n})\right)$$

for every $n \in \mathbb{N}$. We will now argue by induction on $\sum_{i=1}^k a_i$. The start of the induction is the case where $\mathcal{F}$ is a coherent sheaf on one of the surfaces B_i . Since B_i is Cohen-Macaulay, Serre duality gives

$$H^2(B_i, \mathcal{F} \otimes \mathcal{O}_{B_i}(L^{\otimes n})) \simeq \text{Hom}(\mathcal{F} \otimes \mathcal{O}_{B_i}(L^{\otimes n}), \omega_{B_i}).$$

Let $\nu : \tilde{B}_i \rightarrow B_i$ denote the normalisation. Since $\text{Hom}(\mathcal{F}, \omega_{B_i})$ is torsion free, the natural map

$$\text{Hom}(\mathcal{F}, \omega_{B_i}) \rightarrow \nu_*(\nu^*(\text{Hom}(\mathcal{F}, \omega_{B_i})))$$

is injective. Since $\nu_*(\nu^*(\text{Hom}(\mathcal{F}, \omega_{B_i})))$ and $\nu_*(\nu^*(\text{Hom}(\mathcal{F}, \omega_{B_i})/Tor))$ coincide at the generic point of B_i and $\text{Hom}(\mathcal{F}, \omega_{B_i})$ is torsion free, the map

$$\text{Hom}(\mathcal{F}, \omega_{B_i}) \rightarrow \nu_*(\nu^*(\text{Hom}(\mathcal{F}, \omega_{B_i})/Tor))$$

is injective, too. Thus it suffices to show that for any torsion-free sheaf $\mathcal{S}$ on $\tilde{B}_i$ one has

$$H^0(\tilde{B}_i, \mathcal{S} \otimes \mathcal{O}_{\tilde{B}_i}(\tilde{L}^{-\otimes n})) = 0$$

for $n \gg 0$ and $\tilde{L} := \nu^*(L|_{B_i})$. Passing to a desingularisation, we may assume $\tilde{B}_i$ smooth and $\mathcal{S}$ locally free. Fix now a Kähler form ω on $\tilde{B}_i$, and let $\mathcal{G}_1 \subset \mathcal{S}$ be the first sheaf of the Harder-Narasimhan filtration with respect to ω . Since $L|_{B_i}$ is a non-zero nef divisor we have $\tilde{L} \cdot \omega > 0$. Thus there exists a number $n_0 \in \mathbb{N}$ such that $\mathcal{G}_1 \otimes \mathcal{O}_{\tilde{B}_i}(\tilde{L}^{-\otimes n})$ has negative slope for all $n \geq n_0$. In particular $\mathcal{S} \otimes \mathcal{O}_{B_i}(L^{-\otimes n})$ has no global sections.

For the induction step we simply choose a surface B_j such that $a_j > 0$. Then the kernel of the restriction map

$$\mathcal{F} \rightarrow \mathcal{F} \otimes \mathcal{O}_{(\sum_{i=1}^k a_i B_i) - B_j}$$

is a sheaf $\mathcal{G}$ with support on B_j . Thus by the induction hypothesis we know that for $n \gg 0$ the second cohomology vanishes for both $\mathcal{G} \otimes \mathcal{O}_{B_j}(L^{\otimes n})$ and $\mathcal{F} \otimes \mathcal{O}_{(\sum_{i=1}^k a_i B_i) - B_j}(L^{\otimes n})$. $\square$

8.6. Corollary. *Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold, and let L be a nef Cartier divisor on X . Let $D \in |L|$ be effective and set $B := \text{Supp } D$ and $B_1, \dots, B_k$ for the irreducible components of B . Suppose that B_i is Cohen-Macaulay for every $i \in \{1, \dots, k\}$.*

Suppose also that $L|_{B_i} \neq 0$ for every $i \in \{1, \dots, k\}$. Then there exists a number $n_0 \in \mathbb{N}$ and constants $c_1, c_2 \in \mathbb{N}$ such that for all $n \geq n_0$:

$$\dim H^2(X, L^{\otimes n}) = c_1$$

and

$$\dim H^2(X, L^{\otimes n} \otimes \mathcal{O}_X(-B)) = c_2.$$

Proof. For all $n \in \mathbb{N}$ we have an exact sequence

$$0 \rightarrow L^{\otimes n-1} \rightarrow L^{\otimes n} \rightarrow \mathcal{O}_D(L^{\otimes n}) \rightarrow 0.$$

By Lemma 8.5,

$$H^2(D, \mathcal{O}_D(L^{\otimes n})) = 0$$

for all $n \gg 0$. Thus the map $H^2(X, L^{\otimes n-1}) \rightarrow H^2(X, L^{\otimes n})$ is surjective for all $n \gg 0$. This shows the first statement.

For the second statement assume without loss of generality that $D - B \neq 0$. Note that $D - B$ is an effective Weil divisor whose support is contained in B . For all $n \in \mathbb{N}$ we have an exact sequence

$$0 \rightarrow L^{\otimes n-1} \rightarrow L^{\otimes n-1} \otimes \mathcal{O}_X(D - B) \rightarrow \mathcal{O}_{D-B}(L^{\otimes n-1}) \rightarrow 0.$$

Again by Lemma 8.5

$$H^2(D - B, \mathcal{O}_{D-B}(L^{\otimes n-1})) = 0$$

for all $n \gg 0$. Thus the map

$$H^2(X, L^{\otimes n-1}) \rightarrow H^2(X, L^{\otimes n-1} \otimes \mathcal{O}_X(D - B))$$

is surjective for $n \gg 0$ and by the first statement $H^2(X, L^{\otimes n-1})$ is constant for $n \gg 0$. We conclude by noting that

$$L^{\otimes n-1} \otimes \mathcal{O}_X(D - B) \simeq L^{\otimes n} \otimes \mathcal{O}_X(-B)$$

since both sheaves are reflexive and coincide on the smooth locus of X . $\square$

Proof of Theorem 8.2. By Lemma 6.1 and Lemma 6.9 we are reduced to proving the following statement:

Let X be a normal $\mathbb{Q}$ -factorial compact Kähler threefold such that $(X, 0)$ is klt, carrying a divisor $D \in |mK_X|$ with the following properties:

- a) Set $B := \text{Supp } D$. The pair (X, B) is lc and $X \setminus B$ has terminal singularities.*
- b) The divisor $K_X + B$ is nef and we have $\nu(K_X + B) = 2$. Moreover we have $\kappa(X) = \kappa(K_X + B)$.*
- c) For every irreducible component $T \subset B$ we have $(K_X + B)|_T \neq 0$.*
- d) We have $(K_X + B) \cdot K_X^2 \geq 0$.*

Then $\kappa(X) \geq 1$.

Step 1: Singularities of X .

We claim that there is a finite set $S \subset X$ such that $X \setminus S$ has canonical singularities. By hypothesis $X \setminus B$ has only terminal singularities, which are isolated. Thus it remains to consider the singular points of X which are contained in B . Taking

a finite covering of X by analytic neighbourhoods we see that it is sufficient to prove the claim for X a Stein variety. Thus we can take a hyperplane section H of X . Now for general H , the pair (H, B_H) is lc by [Kol97, Prop.7.7], so by [Kaw88, Thm.9.6] every point $p \in B_H \subset H$ is a rational double point in H . Hence by [KM98, Thm.5.34], the point p is a canonical singularity of X .

Step 2: A Chern class inequality.

Let $\mu : X' \rightarrow X$ be a terminal modification of X (cf. Theorem 2.3). Thus X' has only terminal singularities, and there exists an effective $\mathbb{Q}$ -divisor Δ such that

$$K_{X'} + \Delta \sim_{\mathbb{Q}} \mu^* K_X.$$

Let m be the Cartier index of $K_X + B$, then

$$L := m(K_X + B)$$

is Cartier, and we set $L' := \mu^*(L)$. We prove the basic Chern class inequality

$$(36) \quad L' \cdot (K_{X'}^2 + c_2(X')) \geq 0.$$

In fact, since X has only finitely many non-canonical points, $\mu(\Delta)$ is finite. Therefore the projection formula and our assumption d) yield

$$(37) \quad L' \cdot K_{X'}^2 = \mu^* L \cdot K_{X'}^2 = m(K_X + B) \cdot K_X^2 \geq 0.$$

By Proposition 8.4 the sheaf $\Omega_{X'}$ is generically nef. Since

$$K_{X'} + \Delta + \mu^* B \sim_{\mathbb{Q}} \mu^*(K_X + B)$$

is nef, the conditions of Theorem 7.11 are satisfied for $\mathcal{F} := (\Omega_X)^{**}$ and $P := \Delta + \mu^* B$. Having in mind that $L^3 = m^3(K_X + B)^3 = 0$ and using (37), Theorem 7.11 yields

$$L' \cdot c_2(X') \geq 0,$$

hence the Chern class inequality (36) is established.

Step 3: A Riemann-Roch computation. Since $K_X + B$ is nef and $\mathbb{Q}$ -linearly equivalent to an effective divisor with support B , the equality $(K_X + B)^3 = 0$ implies that

$$(38) \quad (K_X + B)^2 \cdot T = 0$$

for every irreducible component $T \subset B$. Since K_X is $\mathbb{Q}$ -linearly equivalent to an effective divisor with support B , we conclude

$$(K_X + B)^2 \cdot K_X = 0.$$

Since $\mu(\Delta)$ is finite the projection formula yields

$$K_{X'} \cdot (L')^2 = K_X \cdot L^2 - \Delta \cdot (\mu^* L)^2 = m^2 K_X \cdot (K_X + B)^2 = 0.$$

Thus Proposition 8.3 gives

$$\chi(X', \mathcal{O}_{X'}(nL')) = nL' \cdot \frac{K_{X'}^2 + c_2(X')}{12} + \chi(X', \mathcal{O}_{X'})$$

for all $n \in \mathbb{N}$. Thus (36) yields a constant k such that

$$\chi(X', \mathcal{O}_{X'}(nL')) \geq k$$

for all $n \in \mathbb{N}$. Since X has rational singularities, we conclude that

$$\chi(X, \mathcal{O}_X(nL)) \geq k$$

for all $n \in \mathbb{N}$. Since $H^3(X, nL) = 0$ for $n \gg 0$ and $\dim H^2(X, nL)$ is constant for $n \gg 0$ by Corollary 8.6 (note that the components of B are Cohen-Macaulay by [KM98, 5.25]), we arrive at

$$(39) \quad h^0(X, nL) \geq h^1(X, nL) + c$$

with some constant $c \in \mathbb{Z}$.

Step 4. A simple case. Although not really necessary, it is instructive to give the simple concluding argument in the case of strict inequality in (36). Then the preceding computation yields that

$$h^0(X, nL) \geq h^1(X, nL) + Dn$$

with some positive constant D . Thus

$$\kappa(K_X + B) = \kappa(X, L) > 0,$$

so $\kappa(X) \geq 1$ by assumption b).

Step 5: Conclusion. By (39) it suffices to show that $h^1(X, nL)$ grows at least linearly. We consider the exact sequence

$$(40) \quad 0 \rightarrow \mathcal{O}_X(nL - B) \rightarrow \mathcal{O}_X(nL) \rightarrow \mathcal{O}_B(nL) \rightarrow 0.$$

By Corollary 8.6 we know that $h^2(X, \mathcal{O}_X(nL - B))$ is constant for $n \gg 0$. Taking cohomology of the exact sequence (40), it remains to show that $h^1(B, \mathcal{O}_B(nL))$ grows at least linearly. To this extent, we will prove that $\chi(B, \mathcal{O}_B(nL))$ is constant. Assuming this for the time being, let us see how to conclude: by Lemma 8.5 we have $H^2(B, \mathcal{O}_B(nL)) = 0$ for $n \gg 0$. Moreover by adjunction [Kwc92, 16.9.1] $\mathcal{O}_B(K_B) \simeq \mathcal{O}_B(K_X + B)$, so by Proposition 5.3

$$h^0(B, \mathcal{O}_B(nL)) = h^0(B, \mathcal{O}_B(nm(K_X + B))) = h^0(B, \mathcal{O}_B(nmK_B))$$

grows linearly. Thus $h^1(B, \mathcal{O}_B(nL))$ grows linearly.

Proof of the claim. By [LR13, Thm.3.1] the Euler characteristic $\chi(B, \mathcal{O}_B(nL))$ on the slc surface B is computed by the usual Riemann-Roch formula¹²

$$\chi(B, \mathcal{O}_B(nL)) = \chi(B, \mathcal{O}_B) + \frac{1}{2}(nL|_B) \cdot (nL|_B - K_B).$$

Yet by (38) we have $0 = L^2 \cdot B = (L|_B)^2$. Since $\mathcal{O}_B(K_B) \simeq \mathcal{O}_B(K_X + B)$ is a multiple of $L|_B$, this also implies that $L|_B \cdot K_B = 0$. $\square$

8.C. Proof of Theorem 1.1.

Proof. By [DP03, Thm.0.3] we have $\kappa(X) \geq 0$. If $\nu(X) = 3$, then K_X is big, hence X is Moishezon and therefore projective [Nam02]. Thus the result follows from the base point free theorem. If $\kappa(X) = \nu(X) \leq 2$ the statement follows from Kawamata's theorem [Kaw85, Thm.1.1], [Nak87, Thm.5.5], [Fuj11b, Sect.4].

By Theorem 8.1 and Theorem 8.2 we are thus left to exclude the possibility that $\kappa(X) = 1$ and $\nu(X) = 2$. This is done exactly as in [Kaw85, Thm.7.3]. $\square$

9. APPLICATIONS

In this concluding section we apply the MMP to explore the structure of non-algebraic compact Kähler threefolds X with Kodaira dimension $\kappa(X) \leq 0$.

¹²We refer to [LR13, Ch.2.3] for the definition of the intersection product on the non-normal surface B .

9.A. Uniruled threefolds.

9.1. Theorem. *Let X be a smooth non-algebraic compact Kähler threefold with $\kappa(X) = -\infty$. Then X is bimeromorphic to a normal compact Kähler threefold X' with at most terminal singularities with the following properties. There exists a contraction*

$$\varphi : X' \rightarrow Y$$

of an extremal ray in $\overline{NA}(X')$ such that

- (1) *Y is a normal non-algebraic Kähler surface with only rational singularities.*
- (2) *There is a finite set $A \subset Y$ such that $\varphi|_{X' \setminus \varphi^{-1}(A)} : X' \setminus \varphi^{-1}(A) \rightarrow Y \setminus A$ is a conic bundle.*
- (3) *φ realises the MRC fibration of X' , and $\kappa(\hat{Y}) \in \{0, 1\}$ for any desingularisation $\hat{Y} \rightarrow Y$.*

Proof. By [HP13a, Thm.1.4] the manifold X is uniruled. We first claim that “the” base B , chosen smooth, of the MRC-fibration $X \dashrightarrow B$ has dimension two. Indeed, if $\dim B = 1$, then the MRC fibration is realised by a morphism $X \rightarrow B$ and the general fibre F is rationally connected, hence $H^2(F, \mathcal{O}_F) = 0$. This immediately implies $H^2(X, \mathcal{O}_X) = 0$, so X is projective by Kodaira’s criterion. The same of course applies when $\dim B = 0$ which is to say that X is rationally connected. Notice also that B is non-algebraic, otherwise X were algebraic.

By [HP13b, Thm.1.1] we can run a MMP which terminates with a Mori fibre space $\varphi : X' \rightarrow Y$. Moreover the properties (1) and (2) are shown in [HP13b, Thm.1.1] and [HP13b, Rem.4.2]. We have seen above that the base of the MRC fibration has dimension two, so φ realises this fibration for X' . In particular Y is not algebraic, so any desingularisation $\hat{Y}$ is not uniruled. Thus we have $\kappa(\hat{Y}) \geq 0$. The non-algebraicity of Y also yields $\kappa(\hat{Y}) \leq 1$. $\square$

We describe the structure of X resp. X' more closely. If the algebraic dimension $a(Y) = 1$, we denote by $f : Y \rightarrow C$ the algebraic reduction.

9.2. Corollary. *In the setting of Theorem 9.1, the algebraic dimensions $(a(X), a(Y))$ can take the following values.*

- (1) $(0, 0)$, and Y is bimeromorphic to a K3-surface or a torus.
- (2) $(1, 0)$, and X is bimeromorphic (up to an étale quotient possibly) of a product of a K3-surface or a tori with an elliptic curve.
- (3) $(1, 1)$, and $f \circ \varphi$ is the algebraic reduction of X' ; the general fiber $f \circ \varphi$ being a ruled surface $\mathbb{P}(V)$ over a (possibly varying) elliptic curve E with $V = \mathcal{O}_E \oplus L$ and $L \equiv 0$ but not torsion or V the non-split extension of two trivial line bundles.
- (4) $(2, 1)$, and $X' \simeq Y \times \mathbb{P}_1$, possibly after a base change $\tilde{C} \rightarrow C$.

Proof. Assertion (1), (2) and (3) are contained in [Fuj87, 14.1], [CP00, 9.1]. As to (4), since it is shown in [CP00, 9.1] that a more precise structure property holds: after passing to $\tilde{X}$ and $\tilde{Y}$ via a finite (ramified) cover $\tilde{C} \rightarrow C$, we obtain

$$\tilde{X} \simeq Z \times_{\tilde{C}} \tilde{Y},$$

where $Z \rightarrow \tilde{C}$ is birational to $\mathbb{P}_1 \times \tilde{C}$, hence the claim. $\square$

9.3. Example. The cases (2) and (4) can obviously be realised by products, we give examples for the other cases.

a) Let Y be a K3 surface with $\text{Pic}(Y) = 0$ and set $X = \mathbb{P}(T_Y)$. Then X does not contain any divisor, hence $a(X) = a(Y) = 0$.

b) Let Y be a two-dimensional torus of algebraic dimension 1, and denote by F the general fibre of the algebraic reduction $f : Y \rightarrow C$. Let $\mathcal{L} \in \text{Pic}(Y)$ be a line bundle such that $\mathcal{L}|_F$ is numerically trivial but not torsion, and set $X := \mathbb{P}(\mathcal{O}_Y \oplus \mathcal{L})$. We claim that $a(X) = a(Y) = 1$: it suffices to prove that $\kappa(X, \mathcal{G}) \leq 1$ for any line bundle $\mathcal{G}$ on X . Now any line bundle $\mathcal{G}$ on X is of the form

$$\mathcal{G} = \mathcal{O}_{\mathbb{P}(\mathcal{O}_Y \oplus \mathcal{L})}(k) \otimes \pi^*(\mathcal{F})$$

with a line bundle $\mathcal{F}$ on Y . If $\kappa(\mathcal{G}) \geq 1$, then $k \geq 0$ and

$$h^0(X, \mathcal{G}^{\otimes m}) = h^0(Y, S^{km}(\mathcal{O}_Y \oplus \mathcal{L}) \otimes \mathcal{F}^m) = \sum_{l=0}^{km} h^0(Y, \mathcal{L}^l \otimes \mathcal{F}^m).$$

Since f is the algebraic reduction, the line bundle $\mathcal{F}$ has degree 0 on F . Since $\mathcal{L}|_F$ is not torsion there exists a unique $l_0 \in \{0, \dots, km\}$ such that $(\mathcal{L}^l \otimes \mathcal{F}^m)|_F \simeq \mathcal{O}_F$. Thus we have

$$h^0(X, \mathcal{G}^{\otimes m}) = h^0(Y, \mathcal{L}^{l_0} \otimes \mathcal{F}^m),$$

We immediately obtain $\kappa(X, \mathcal{G}) \leq 1$.

9.B. Threefolds with trivial canonical bundles. We will study the Albanese map for certain threefolds with terminal singularities (cf. [BS95, 2.4.1] for the existence of Albanese maps in the presence of rational singularities).

9.4. Lemma. *Let X be a (non-algebraic) compact Kähler threefold with terminal singularities. If $\kappa(X) = 0$, then the Albanese map $\alpha : X \rightarrow A = \text{Alb}(X)$ is surjective with connected fibres. In particular we have $q(X) \leq 3$.*

Proof. Apply [Uen87, Main Thm.I] to a desingularisation $\hat{X} \rightarrow X$. □

9.5. Theorem. *Let X be a non-algebraic compact Kähler threefold with terminal singularities. If $K_X \equiv 0$, there exists a Galois cover $f : \tilde{X} \rightarrow X$ that is étale in codimension one such that either $\tilde{X}$ is a torus or a product of an elliptic curve and a K3 surface.*

Proof. By Theorem 1.1, the divisor K_X is semi-ample. Thus we can choose $m \in \mathbb{N}$ minimal such that $\mathcal{O}_X(mK_X) = \mathcal{O}_X$. Let $X' \rightarrow X$ be the induced cyclic cover, then we have $\mathcal{O}_{X'}(K_{X'}) \simeq \mathcal{O}_{X'}$. In particular X' is Gorenstein with terminal singularities [KM98, Cor.5.21]. Thus the Riemann-Roch formula 8.3 gives $\chi(X', \mathcal{O}_{X'}) = 0$. Since $h^2(X', \mathcal{O}_{X'}) \geq 1$, the variety X' being non-algebraic with rational singularities only, we conclude that $h^1(X', \mathcal{O}_{X'}) \geq 1$.

Now consider the Albanese map $\alpha : X' \rightarrow \text{Alb}(X') =: A$. By Lemma 9.4 the morphism α is surjective with connected fibres.

Case 1: $q(X') = 3$. Then α is bimeromorphic. Since A is smooth, $K_{X'} \equiv E$ with E an effective divisor whose support equals the α -exceptional locus. Since $K_{X'} \equiv 0$ we conclude that $X' \simeq A$.

Case 2: $q(X') = 2$. By [Uen87, Main Thm.I,2)] there exists a finite set $S \subset A$ such that α is an elliptic bundle over $A \setminus S$. By [CP00, Prop.6.7(i)] this implies

that (after finite étale base change) the fibration α is bimeromorphic to a compact Kähler manifold X'' that is an elliptic bundle $X'' \rightarrow A$. Yet a Kähler manifold which is an elliptic bundle over a torus is an étale quotient of a torus [Bea83], so after finite étale cover we have $q(X') = q(X'') = 3$. We conclude by applying Case 1.

Case 3: $q(X') = 1$. By [Uen87, Main Thm.I,3)] there exists an analytic fibre bundle $X^* \rightarrow A$ that is bimeromorphic to X' . Since the Kodaira dimension is an invariant of varieties with terminal singularities, we have $\kappa(X^*) = 0$. In particular the general fibre F of $X^* \rightarrow A$ is the blow-up of a torus or a K3 surface¹³. We next run a relative MMP over the elliptic curve A . Since $X^* \rightarrow A$ is a fibre bundle, every step of this MMP is the blow-up along an étale multisection of the fibration, so the outcome is an analytic fibre bundle $\bar{X} \rightarrow A$ such that the general fibre F is a torus or K3 surface. If F is covered by a torus, then $\bar{X}$ is a torus after finite étale cover, see e.g. [Bea83]. If F is a K3 surface, the fibre bundle trivialises after finite étale base change $A' \rightarrow A$ ([Fuj78, Cor.4.10], cf. [CH12, Lemma 2.15] for more details). Thus we have $\bar{X} \simeq A' \times F$. In conclusion we see that (up to finite étale cover) we have a bimeromorphic map

$$\mu : X' \dashrightarrow \bar{X}$$

with $\bar{X}$ a torus or a product $A' \times F$. Since both $K_{X'}$ and $K_{\bar{X}}$ are numerically trivial, we see that μ is an isomorphism in codimension one [Han87], [Kol89, 4.3]. Moreover μ decomposes into a finite sequence of flops by [Kol89, 4.9]. Note however that the last flop of this sequence yields a rational curve in $\bar{X}$ that is very rigid (in the sense of [HP13a, Defn.4.3]). Since $\bar{X}$ is a torus or a product $A' \times F$ such a curve does not exist on $\bar{X}$. Thus μ is an isomorphism. $\square$

Using the existence of minimal models for a smooth compact Kähler threefold, we deduce

9.6. Corollary. *Let X be a non-algebraic compact Kähler threefold with $\kappa(X) = 0$. There exists a finite cover which is bimeromorphic to a torus or a product of an elliptic curve and a K3 surface.*

We can now derive Theorem 1.2 from Theorem 9.5:

Proof of Theorem 1.2. Since X is not uniruled, K_X is pseudo-effective [Bru06]. By [HP13a, Thm.1.1] there exists a minimal model $X \dashrightarrow X'$. Since X has algebraic dimension zero, we see that $\kappa(X) = 0$. Thus we have $\nu(X') = \kappa(X') = 0$ by Theorem 1.1, i.e. the canonical divisor $K_{X'}$ is numerically trivial. Since X (and hence X') is not covered by curves Theorem 9.5 yields that $X' \simeq T/G$ with T a torus and G a finite group. Since X (and hence X') is not covered by positive-dimensional subvarieties, the torus T has no positive-dimensional subvarieties. In particular T/G has no positive-dimensional subvarieties, so $X \dashrightarrow T/G$ extends to a morphism. $\square$

¹³If F was bimeromorphic to an Enriques or bielliptic surface we would have $0 = H^2(X^*, \mathcal{O}_{X^*}) = H^2(X', \mathcal{O}_{X'})$, a contradiction.

REFERENCES

- [Anc87] Vincenzo Ancona. Vanishing and nonvanishing theorems for numerically effective line bundles on complex spaces. *Ann. Mat. Pura Appl. (4)*, 149:153–164, 1987.
- [Ara10] Carolina Araujo. The cone of pseudo-effective divisors of log varieties after Batyrev. *Math. Z.*, 264(1):179–193, 2010.
- [BCE⁺02] Thomas Bauer, Frédéric Campana, Thomas Eckl, Stefan Kebekus, Thomas Peterzell, Sławomir Rams, Tomasz Szemberg, and Lorenz Wotzlaw. A reduction map for nef line bundles. In *Complex geometry (Göttingen, 2000)*, pages 27–36. Springer, Berlin, 2002.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010.
- [BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *Journal of Algebraic Geometry*, 22:201–248, 2013.
- [Bea83] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18(4):755–782 (1984), 1983.
- [BHPVdV04] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, second edition, 2004.
- [Bou04] Sébastien Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004.
- [BPEG13] Sébastien Boucksom; Philippe Eyssidieux and Vincent Guedj, editors. *An introduction to the Kähler-Ricci flow. Selected papers based on the presentations at several meetings of the ANR project MACK*. Cham: Springer, 2013.
- [Bru06] Marco Brunella. A positivity property for foliations on compact Kähler manifolds. *Internat. J. Math.*, 17(1):35–43, 2006.
- [BS94] Shigetoshi Bando and Yum-Tong Siu. Stable sheaves and Einstein-Hermitian metrics. In *Geometry and analysis on complex manifolds*, pages 39–50. World Sci. Publ., River Edge, NJ, 1994.
- [BS95] Mauro C. Beltrametti and Andrew J. Sommese. *The adjunction theory of complex projective varieties*, volume 16 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1995.
- [Cao13] Junyan Cao. A remark on compact Kähler manifolds with nef anticanonical bundles and its applications. *arXiv preprint*, 1305.4397, 2013.
- [CH12] Benoît Claudon and Andreas Höring. Compact Kähler manifolds with compactifiable universal cover. *arXiv preprint*, 2012.
- [Cor07] Alessio Corti, editor. *Flips for 3-folds and 4-folds*, volume 35 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2007.
- [CP00] Frédéric Campana and Thomas Peternell. Complex threefolds with non-trivial holomorphic 2-forms. *J. Algebraic Geom.*, 9(2):223–264, 2000.
- [CT13] Tristan C. Collins and Valentino Tosatti. Kähler currents and null loci. *arXiv preprint*, 1304.5216, 2013.
- [Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [DP03] Jean-Pierre Demailly and Thomas Peternell. A Kawamata-Viehweg vanishing theorem on compact Kähler manifolds. *J. Differential Geom.*, 63(2):231–277, 2003.
- [DP04] Jean-Pierre Demailly and Mihai Paun. Numerical characterization of the Kähler cone of a compact Kähler manifold. *Ann. of Math. (2)*, 159(3):1247–1274, 2004.
- [Eno88] Ichiro Enoki. Stability and negativity for tangent sheaves of minimal Kähler spaces. In *Geometry and analysis on manifolds (Katata/Kyoto, 1987)*, volume 1339 of *Lecture Notes in Math.*, pages 118–126. Springer, Berlin, 1988.
- [Fuj78] T. Fujita. On Kähler fiber spaces over curves. *J. Math. Soc. Japan*, pages 779–794, 1978.
- [Fuj87] Takao Fujita. On polarized manifolds whose adjoint bundles are not semipositive. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 167–178. North-Holland, Amsterdam, 1987.

- [Fuj07] Osamu Fujino. Special termination and reduction to pl flips. In *Flips for 3-folds and 4-folds*, volume 35 of *Oxford Lecture Ser. Math. Appl.*, pages 63–75. Oxford Univ. Press, Oxford, 2007.
- [Fuj11a] Osamu Fujino. Non-vanishing theorem for log canonical pairs. *J. Algebraic Geom.*, 20(4):771–783, 2011.
- [Fuj11b] Osamu Fujino. On Kawamata’s theorem. In *Classification of algebraic varieties*, EMS Ser. Congr. Rep., pages 305–315. Eur. Math. Soc., Zürich, 2011.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, second edition, 1998.
- [Gri10] Julien Grivaux. Chern classes in Deligne cohomology for coherent analytic sheaves. *Math. Ann.*, 347(2):249–284, 2010.
- [Han87] Masaki Hanamura. On the birational automorphism groups of algebraic varieties. *Compositio Math.*, 63(1):123–142, 1987.
- [HP13a] Andreas Höring and Thomas Peternell. Minimal models for Kähler threefolds. *arXiv preprint*, 1304.4013, 2013.
- [HP13b] Andreas Höring and Thomas Peternell. Mori fibre spaces for Kähler threefolds. *arXiv preprint*, 1310.5837, 2013.
- [Kaw85] Y. Kawamata. Pluricanonical systems on minimal algebraic varieties. *Invent. Math.*, 79(3):567–588, 1985.
- [Kaw88] Yujiro Kawamata. Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces. *Ann. of Math. (2)*, 127(1):93–163, 1988.
- [Kaw91] Yujiro Kawamata. On the length of an extremal rational curve. *Invent. Math.*, 105(3):609–611, 1991.
- [Kaw92a] Yujiro Kawamata. Abundance theorem for minimal threefolds. *Invent. Math.*, 108(2):229–246, 1992.
- [Kaw92b] Yujiro Kawamata. Termination of log flips for algebraic 3-folds. *Internat. J. Math.*, 3(5):653–659, 1992.
- [KK10] János Kollár and Sándor J. Kovács. Log canonical singularities are Du Bois. *J. Amer. Math. Soc.*, 23(3):791–813, 2010.
- [KM92] János Kollár and Shigefumi Mori. Classification of three-dimensional flips. *J. Amer. Math. Soc.*, 5(3):533–703, 1992.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti.
- [Kob87] Shoshichi Kobayashi. *Differential geometry of complex vector bundles*, volume 15 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1987. Kanô Memorial Lectures, 5.
- [Kol89] János Kollár. Flops. *Nagoya Math. J.*, 113:15–36, 1989.
- [Kol97] János Kollár. Singularities of pairs. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 221–287. Amer. Math. Soc., Providence, RI, 1997.
- [Kol07] János Kollár. Kodaira’s canonical bundle formula and adjunction. In *Flips for 3-folds and 4-folds*, volume 35 of *Oxford Lecture Ser. Math. Appl.*, pages 134–162. Oxford Univ. Press, Oxford, 2007.
- [Kol13] János Kollár. *Singularities of the minimal model program*, volume 200 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013. With a collaboration of Sándor Kovács.
- [Kwc92] János Kollár (with 14 coauthors). *Flips and abundance for algebraic threefolds*. Société Mathématique de France, Paris, 1992. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991, Astérisque No. 211 (1992).
- [LR13] Wenfei Liu and Sönke Rollenske. Geography of Gorenstein stable log surfaces. *arXiv preprint*, 1307.1999, 2013.
- [Miy87] Yoichi Miyaoka. The Chern classes and Kodaira dimension of a minimal variety. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 449–476. North-Holland, Amsterdam, 1987.
- [Miy88] Yoichi Miyaoka. Abundance conjecture for 3-folds: case $\nu = 1$. *Compositio Math.*, 68(2):203–220, 1988.

- [Mor88] Shigefumi Mori. Flip theorem and the existence of minimal models for 3-folds. *J. Amer. Math. Soc.*, 1(1):117–253, 1988.
- [Nak87] Noboru Nakayama. The lower semicontinuity of the plurigenera of complex varieties. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 551–590. North-Holland, Amsterdam, 1987.
- [Nam02] Yoshinori Namikawa. Projectivity criterion of Moishezon spaces and density of projective symplectic varieties. *Internat. J. Math.*, 13(2):125–135, 2002.
- [Pau98] Mihai Paun. Sur l’effectivité numérique des images inverses de fibrés en droites. *Math. Ann.*, 310(3):411–421, 1998.
- [Pet98] Thomas Peternell. Towards a Mori theory on compact Kähler threefolds. II. *Math. Ann.*, 311(4):729–764, 1998.
- [Pet01] Thomas Peternell. Towards a Mori theory on compact Kähler threefolds. III. *Bull. Soc. Math. France*, 129(3):339–356, 2001.
- [Sak84] Fumio Sakai. Weil divisors on normal surfaces. *Duke Math. J.*, 51(4):877–887, 1984.
- [Sho92] V.V. Shokurov. 3-fold log flips. Appendix by Yujiro Kawamata: The minimal discrepancy coefficients of terminal singularities in dimension three. *Russ. Acad. Sci., Izv., Math.*, 40(1):95–202, 1992.
- [TT86] Domingo Toledo and Yue Lin L. Tong. Green’s theory of Chern classes and the Riemann-Roch formula. In *The Lefschetz centennial conference, Part I (Mexico City, 1984)*, volume 58 of *Contemp. Math.*, pages 261–275. Amer. Math. Soc., Providence, RI, 1986.
- [Uen87] Kenji Ueno. On compact analytic threefolds with nontrivial Albanese tori. *Math. Ann.*, 278(1-4):41–70, 1987.
- [Var89] Jean Varouchas. Kähler spaces and proper open morphisms. *Math. Ann.*, 283(1):13–52, 1989.
- [Voi02] Claire Voisin. *Théorie de Hodge et géométrie algébrique complexe*, volume 10 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2002.

FRÉDÉRIC CAMPANA, INSTITUT ELIE CARTAN, UNIVERSITÉ HENRI POINCARÉ, BP 239, F-54506. VANDOEUVRE-LES-NANCY CÉDEX, ET: INSTITUT UNIVERSITAIRE DE FRANCE,
E-mail address: frederic.campana@univ-lorraine.fr

ANDREAS HÖRING, LABORATOIRE DE MATHÉMATIQUES J.A. DIEUDONNÉ, UMR 7351 CNRS, UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS, 06108 NICE CEDEX 02, FRANCE
E-mail address: hoering@unice.fr

THOMAS PETERNELL, MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY
E-mail address: thomas.peternell@uni-bayreuth.de